

*Schizura concinna* S. & A.

*Schizura semirufescens* Walk., probably var. *perangulata* Hy Edw.

*Schizura unicornis* S. & A.

*Schizura badia* Pack.

*Schizura leptinoides* Grt.

*Cerura scitcripta* Walk., var. *multiscripta* Riley.

*Cerura occidentalis* Lint.

*Harpyia cinerea* Walk.

*Harpyia scolopendrina* Bdv. As figured by Holland.

*Gluphisia septentrionalis* Walk.

*Gluphisia* sp. Two males, one dated June 3rd, 1905, apparently distinct from anything else known to me, but near *septentrionalis*. I compared a Winnipeg specimen with Packard's figures some years ago, and noted that it resembled *wrightii* as there figured, but it cannot be *wrightii*. I submitted a specimen to Messrs. Barnes and McDunnough, who said it was "nearest *lintneri* which we have from the same general region." It certainly is not *lintneri*.

#### Liparidæ.

*Hemerocampa leucostigma* S. & A.

*Hemerocampa definita* Pack. A series, agreeing with Holland's figure.

*Alone vagans* B. & McD., var. *grisea* B. & McD. This is the species which has always been widely known as *plagiata* Walk., but that species is the same as *definita* Pack. The form is normally brown or grey-brown, but some of the Heath specimens were very blackish grey.

#### Lasiocampidæ.

*Malacosoma fragilis* Stretch. The species was so named in the collection. I am not sure that *americana*, *fragilis* and *pluvialis* are always separable except by larval characters.

#### Platypterygidæ.

*Oreta rosea* Walk.

*Drepana arcuata* Walk.

*Falcaria bilineata* Walk.