middle and high latitude regions of the northern hemisphere. Since the end of the last ice age, about 10,000 years ago, global surface temperatures have probably fluctuated by little more than 1°C. Some fluctuations have lasted several centuries, including the Little Ice Age which ended in the nineteenth century and which appears to have been global in extent.

The changes predicted to occur by about the middle of the next century due to increases in greenhouse gas concentrations from the Business-as-Usual emissions will make global mean temperatures higher than they have been in the last 150,000 years.

The rate of change of global temperatures predicted for Business-as-Usual emissions will be greater than those which have occured naturally on earth over the last 10,000 years, and the rise in sea level will be about three to six times faster than that seen over the last 100 years or so.

Has man already begun to change the global climate?


The instrumental record of surface temperature is fragmentary until the midnineteenth century, after which it slowly improves. Because of different methods of measurement, historical records have to be harmonised with modern observations, introducing some uncertainty. Despite these problems we believe that a real warming of the globe of 0.3°C - 0.6°C has taken place

over the last century; any bias due to urbanisation is likely to be less than 0.05°C.

Moreover since 1900 similar temperature increases are seen in three independent data sets: one collected over land and two over the oceans. The figure below shows current estimates of smoothed global mean surface temperature over land and ocean since 1860. Confidence in the record has been increased by their similarity to recent satellite measurements of mid-tropospheric temperatures.

Although the overall temperature rise has been broadly similar in both hemispheres, it has not been steady, and differences in their rates of warming have sometimes persisted for decades. Much of the warming since 1900 has been concentrated in two periods, the first between about 1910 and 1940 and the other since 1975; the five warmest years on record have all been in the 1980s. The northern hemisphere cooled between the 1940s and the early 1970s when southern hemisphere temperatures stayed nearly constant. The pattern of global warming since 1975 has been uneven with some regions, mainly in the northern hemisphere, continuing to cool until recently. This regional diversity indicates that future regional temperature changes are likely to differ considerably from a global average.

The conclusion that global temperature has been rising is strongly supported by the retreat of most mountain glaciers of the world since the end of the nineteenth century and the fact that global sea level has risen over the same period by an average of 1 to 2mm per year. Estimates of thermal expansion of the oceans, and of



Global mean combined land-air and sea-surface temperatures, 1861 - 1989, relative to the average for 1951-80.