

*Ochria Buffaloensis*, n. s.

♀. The clypeus has a frontal horn, else the insect looks like *Gortyna rutila* and allies. The primaries are brownish red with the extra-basal and subterminal spaces washed with purple. T. a. line geminate, waved, with rather a deep sinus on vein 1, its inner line purple, its outer dark brown red. Orbicular spherical, yellow white. The accessory spots are totally wanting; this last superficial character will separate the moth from any N. Am. species of *Gortyna* allied to *rutila*, known to me or described by Gueneé. Reniform moderate, yellowish, interlaced with a double brown curved line. T. p. line double, nearly straight, not exerted opposite the cell, but prolonged on costa as in *purpurifascia*; its outer line is dark purplish, more diffuse. S. t. line dark, distinct, irregularly dentate. External margin even, bulging opposite median nervules. Veins obscurely purplish. Hind wings pale red, with a mesial straight dark line. Beneath light purple red, with a distinct common line; on hind wings a narrow lunule. Body beneath concolorous with wings; thorax above darker, more purplish. Expanse 40 mil. Miss Mary Walker, Buffalo.

*Polia pallifera*, n. s.

♀. This species resembles Herrich-Schaeffer's figures of *platinea*. Fore wings whitish gray with the median space washed with olivaceous beyond the olive median shade line. Claviform olivaceous, very large, finely lined with black. Reniform and orbicular gray, shaded with olive, the former white narrowly margined with black. Median lines geminate, of the usual shape. Beyond the t. p. line the wing is whitish gray, cut by the olivaceous shaded s. t. line. Fringes obscure with an interrupted dark line. Hind wings fuscous, paler at base with a sinuate mesial line and a pale subterminal shade. Beneath dirty whitish with double lines and discal marks. Body beneath and abdomen obscure whitish or dusty gray; thorax more purely gray. Expanse 42 mil. Illinois, Mr. Bean.

I cannot identify this with any of Mr. Morrison's descriptions under this genus. It has the fascies of European species of the group. It recalls the Californian *Dian. insolens*, but the eyes are naked.

*Homoptera Woodii*, n. s.

♂. This species is more strigate than any other known to me. Collar brown with a black mesial line, tipped with gray. Wings dentate, covered with dark strigae. Base of primaries blackish, defined by a broad velvety