

AGAINST ERYSIPELAS.

[REV. GEN. DE CLIN.]

PAINT.

Tannic Acid } of each 1 part.
Camphor }
Ether 3 parts

Apply with a brush every two hours.

INFLUENZA TRATMENT.

[HUCHARD—AM. M.-S. BULL.]

PILLS.

Quinine Hydrobromate . . } of each 2 Gms.
Sodium Benzoate } [31 grs.]
Caffeine }
Make into 30 pill.—Two or three daily.

AGAINST THE IRRITATIVE COUGH OF PHTHISIS.

[F. H. HENRY—KANS. CITY. MED. IND.]

MIXTURE.

Potassium Cyanide . . . } of each 1 grain
Morphine Acetate . . . } [6.5.ctg.]
Vinegar Sanguinaria . . 2 fl. drs. [7.5. c. c.]
Syrup Tolu 1 fl. oz. [30 "]
Distilled Water enough to make 3 fl. oz. [89 c. c.]

Teaspoonful every three hours.

CONSTIPATION IN CHILDREN.

I. SIMON.—L'UNION MED.]

MIXTURE.

Tincture Cascarella . . . }
Tincture Rhubarb } of each 2 parts.
Tincture Cinnamon . . . }
Tincture Calumbo }
Tincture Gentian }
Tincture Nux Vomica 1 part.
10-20 drops twice daily.

VULVAR ECZEMA.

[L'UN. MED.]

LOTION.

Potassium Bicarbonate 1 part.
Sodium Bicarbonate } of each 2 parts.
Glycerin }
Tincture Opium }
Distilled 60 "

Apply warm.— At night, dust the parts with the following

Powdered Camphor 1 part.
Powdered Starch 40 parts.

SCROTAL ECZEMA.

CAMPBELL—MED. REC.]

LOTION.

Iodoform 2 drams [12 Gms.].
Zinc Oxide 1½ oz. [46 "]
Spirit Camphor 3 drams [11 c. c.]
Lime Water } of each 1½ fl. oz. [45 "]
Linseed Oil }

Apply at night, and have the patient wear a suspensory bandage.

ICHTHYOL IN ACNE ROSACEA.

[PETRINI—SEM. MED.]

PAINT.

Ichthyol 2 parts.
Resorcin 1 part.
Collodion 30 parts.

Apply for three consecutive days. If there are pustules, open them before using the paint.

SPIRITS OF TURPENTINE IN THE TREATMENT OF BURNS.

H. L. McInnis (New York Medical Record) says that spirits of turpentine applied to a burn of either the first, second or third degree will almost at once relieve the pain. The burn will heal much more rapidly than by any other treatment in the author's experience. He applies the turpentine as follows: After wrapping a thin layer of absorbent cotton over the burn, he saturates it with the common commercial turpentine, which is generally found in every house, and then bandages. Being volatile, the turpentine evaporates, and it is therefore necessary to keep the cotton moistened with it. When there are large blebs, he opens them on the second or third day. It is best to keep the spirits off the healthy skin, if possible, as sometimes pain is produced by its action.—British Medical Journal.

PAINFUL PHARYNGITIS.

[F. INGALS—MED. WEEK.]

PAINT.

Morphine Sulphate . . . 3 grains [20 ctg.]
Carbolic Acid } of each 30 " [2 Gms.]
Tannic Acid }
Glycerin } of each ½ fl. oz. [15 c. c.]
Distilled Water }

Apply to the painful part of the throat.