162. Leptura nigrella Say.

No data.

- 163. Leptura carbonata LeConte. Northern California.
- 164. Leptura subcostata Fall.

One specimen taken by Fall at Tahoe.

165. Leptura dehiscens LeConte.

Rare, in the Sierras; northern, one specimen taken by Van Dyke in Siskiyou County. Four taken from small limbs of *Pinus ponderosa* at Calistoga by author. May 28.

166. Leptura sanguinea LeConte.

Found in Northern California and throughout the Sierras on veratrin and other flowers. May 30-July 17.

167. Leptura lætifica LeConte.

Taken in Sierras and Northern California; common in places in Shasta Lake, and Sonoma Counties on flowers of Eriogonum, etc. May 25-June 19.

168. Leptura pernigra Linnell.

Taken in Sierras Madre Mts., Los Angeles County by Van Dyke.

169. Leptura quadrillum LeConte.

Found on yarrow, etc., in the northern part of the State.

170. Leptura chrysocoma Kirby.

Common in the Sierras and Northern California on veratrin, yarrow, and other flowers. Taken in large numbers at Donner Lake and Tahoe by the author in July. July 1–21.

171. Leptura dolorosa LeConte.

Found especially on the Ceanothus in Northern California and the Sierras: common in places.

172. Leptura crassipes LeConte.

Found throughout the State, common in the northern part, less common in the south. Taken by Fall in the San Bernardino Mts. Usually found on the flowers of the yarrow, ox-eyed daisy, etc. Bred from *Umbellularia californica* by Rivers and Van Dyke. Bred from *Eucalyptus globulus* by Van Dyke. Taken by author at Calistoga. May 21.