

Genus 2. *Melanippe*, Dup.

1. *M. Gothicata*, Guén.—Common, June.

Distribution.—Rouge District, Sorel and Southern shores of the gulf.

2. *M. lacustrata*, Guén.—Abundant, May and August.
3. *M. intermediata*, Guén.—Common, May, July and August.

Genus 3. *Coremia*, Guén.

1. *C. propugnata*, Meir. Verz.—August. (Eu).

Genus 4. *Scotosia*, Steph.

1. *S. undulata*, Linn.—Common, June and July. (Eu).

Distribution.—Massachusetts, (Harris); Rouge District.

2. *S. affirmaria*, Walker, M.S.S. N. sp.—Abundant in August.

"*Mas.* Rufo-cinerea; thorax nigro fasciatus; alæ anticæ lineis plurimis rufescentibus subundulatis, lineis tribus distinctis dentatis nigris, linea submarginali elunulis albidis, litura discali nigra obliqua minima."

"*Male*. Cinereous. Body above and forewings much tinged with red. Thorax with a black band. Wings with a black marginal festoon. Forewings with many slight undulating reddish lines, with three distinct dentated black lines, and with a whitish submarginal line which is composed of lunules; discal mark black, oblique, very small. Hindwings pale cinereous, with several indistinct darker lines and with a reddish tinge about the interior border and the exterior border. Length of the body, 8 lines; of the wings, 18 lines." Walker, M.S.S.

Group 5. PYRALIDINA, Stainton.

Sub. group 1. PYRALIDES, Guén.

Section 1. DELTOIDES, Guén.

Family, HYPENIDÆ, Guén.

Genus 1. *Hypena*, Schr.

1. *H. N. sp.*—Abundant flying by day on the Mountain, May to July.
2. *H. cæcalis*, Walker.—May.

Genus 2. *Hormissa*.

1. *H. effusalis*, Walker, M.S.S.—Common in July.

"*Mas.* Cerrino-cinerea; palpi decumbentes, longissimi articulo 2o. apicem versus arcuato, 3o. lanceolato; antennæ pectinatæ; alæ anticæ lineis tribus fusciscentibus subundulatis, linea marginali fusca tenui, reniformi e striga angusta arcuata; posticæ striga latissima, lineis indistinctis."

"*Male*, Cinereous, with a slight fawn coloured tinge. Palpi decumbent, very long; 2d. joint much curved towards the tip;