
A LIST OF HEMIPTERA HETEROPTERA COLLECTED IN
SOUTH LOUISIANA.

BY C. H. T. TOWNSEND, CONSTANTINE, MICH.

Now that we have been supplied by Mr. Uhler with a list of the Hemiptera Heteroptera of North America, which has been much needed, it will be interesting to know more about the geographical distribution of the species. The following list is short, but will nevertheless add to our knowledge of these insects. The species were collected at the same time and in the same localities with the Coleoptera given in my list (CAN. ENTOM., xvii., p. 66-73), and the preliminary notes there given apply also here. The species were kindly determined for me by Mr. Uhler.

PENTATOMIDÆ.

Mineus bioculatus Fab. One specimen taken on plants.*

Podisus sp. The nymphs of two species of *Podisus* were taken on wild plants; one black and reddish, and three smaller of a greenish color.

Euthyrhynchus floridanus Linn. One taken 25th May on plants along the wild side of a ditch, on the edge of a plantation. B. la F.

Ocbealus typhæus Fab. Two taken on plants. This species is not given in the Check List.

Euschistus sp. Two nymphs were taken on plants; they are probably two different species, though they look much alike.

Proxys punctulatus Pal. Beauv. Twenty taken, nearly all on 29th March, under dry logs near Lake Pontchartrain, at Milneburg.

Nezara viridula Linn. Two imagos taken 2nd June on plants, and one nymph at another time. B. la F.

Nezara hilaris Say. One nymph taken on plants.

Edessa bifida Say. One taken about first of June, I think, on wild plants. B. la F.

COREIDÆ.

Metapodius granulatus Dallas. Numbers taken on thistles and other plants along ditches on plantations, and in other open sunny places. N. O.; B. la F. This is probably not the species so

* Where no locality is given it may be either, but is probably B. la F. Where no date, it is unknown, unless the species was more or less abundant during my stay.