

- than 2 and 3 together, but as long or longer than 3 and 4 together; in middle tarsi joint 5 equals twice 4, equals or is slightly shorter than 2, which is shorter than 1 and shorter than 3 and 4 together; in posterior tarsi joint 2 is less than three times 4, much longer than 5, and nearly as long as 3 and 4 together. F.
- BB. Hind femora without a row of minute bristles on the side; eyes small, near lower edge of head; antennal groove in anterior half of head; maxillary palpi in female with joint 2 more than three-fourths of 4; head in female evenly rounded from occiput to mouth; pronotal comb of 18 spines; colour light brown, sometimes darker dorsally. C.
- C. Abdomen more convex above than below, and without teeth on discs of first two or three segments above; bristles on second antennal joint shorter than third joint; bristles on abdomen as follows: first row on each side of dorsal segments with 4 or 5 bristles, second row with 6, ventral segments with 4 on each side; in anterior tarsi joint 5 is longer than 3 and 4 together; in middle tarsi joint 5 equals twice 4 and equals 1; in posterior tarsi joint 5 equals two-thirds of 2, is less than twice 4, and equals 3, 2 is slightly less than 3 and 4 together, while 1 is more than twice 5; colour light brown; length: female, 3 mm. *Wickhami*, n. sp.
- CC. Abdomen concave or very slightly convex above; with minute teeth on discs of first two or three abdominal segments above. D.
- D. Bristles on second antennal joint nearly as long as third joint; bristles on abdomen as follows: first row on each side of dorsal segments with 6 bristles, second row with 7, ventral segments with 4 or 5 on each side; in anterior tarsi joint 5 nearly as long as 3 and 4 together; in middle tarsi joint 5 is less than 1, about as long as 2 or shorter, and less than 3 and 4 together; in posterior tarsi joint 5 equals two-thirds of 2, equals 3, and is less than twice 4, 2 is less than three times 4, and 1 is more than twice 5; colour light brown; length: female, 2.5-3 mm. *Gillettei*, n. sp.
- DD. Bristles of second antennal joint short; comparative lengths of tarsal joints not as above. E.
- E. Male claspers unarmed; head evenly rounded; first and second rows of bristles on dorsum of abdominal segments each with 7 bristles; in anterior tarsi joint 5 equals 3 and 4 together; in middle tarsi joint 1 equals 2, equals 5, and equals 3 and 4 together;