

- Homalodisca liturata* Ball. Occasional specimens taken in southern counties. Swept from weeds.
- Gypona modesta* Spangb. Two adults captured at Poplar Bluff.
- Gypona bimaculata* Spangb. Collected by sweeping rank growth in marshes near Charleston.
- Platymetopius acutus* var. *dubius* Van D. Rather numerous on grass in southwestern part of the State.
- Deltocephalus melscheimeri* Fh. Occasional specimens captured. Occurs throughout the State.
- Athysanus comma* Van D. One adult taken at St. Louis.
- Athysanus anthracinus* Van D. Rather numerous in western counties.
- Phlepsius fulvidorsum* Fh. Numerous on willows growing along creeks and rivers in Southwestern Missouri.
- Phlepsius nebulosus* Van D. Occasional specimens taken near Charleston.
- Thamnotettix brittoni* Osb. A few adults were swept from weeds at Charleston.
- Chlorotettix balli* Osb. Specimens captured were all from northern parts of State.
- Chlorotettix lusorius* O. & B. Occurs most numerous in eastern counties.
- Jassus melanotus* Spangb. Collected by F. M. Moody at Branson, by sweeping weeds growing on high ridges of the Ozarks.
- Tinobregmus pallidus* Osb. Two adults captured by F. M. Moody from low shrubs growing wild in the woods at Branson.
- Cicadula punctifrons* Fall. Few adults captured at Dexter.
- Empoasca flavescens* Fabr. Abundant on willows at Branson.
- Empoasca trifasciata* Gill. One specimen captured at Charleston, July 26.
- Erythroneura illinoiensis* Gill. Rather abundant in eastern counties.
- Erythroneura crevecœuri* Gill. Not common. Occasional specimens taken at Charleston.