

- Legs with the femora dark brown at the tip with an indistinct sub-apical ring; a cross-vein in cell R_3 ; seam on cord pale brown, broad; antennae pale.....*translucida*, sp. n.
(Panama)
5. Wings with an abundant pattern in the cells.....*gloriosa*, sp. n.
(Guatemala)
- Wings with the markings scanty and more or less confined to the neighbourhood of veins.....6.
6. Wing hyaline, with the markings brown; pleurae with a brown band; tibiae and tarsi uniform dark.....*eiseni*, sp. n.
(Guatemala)
- Wing dusky, with the markings dark brown; no pleural band; tibiae at tip, and tarsi, orange brown.....*lutsi*, sp. n.
(Brit. Guiana)

Furcomyia reticulata, sp. n.

Antennae brown; thorax yellow, with an irregular brown median stripe; legs yellow, darkening to brown apically; wings hyaline, reticulated with brown marks.

♀.—Length, 4.5–6 mm; wing, 5.3–5.4 mm.

♀.—Head: rostrum yellowish brown; palpi dark brownish black. Antennae, basal segments pale, whitish; flagellum light brown, the segments rounded, becoming oval and then elongated toward the tip of the antennae. Front, vertex and occiput dull yellow, the vertex and occiput prolonged caudad, with two brown stripes above and brown on the sides.

Thorax: pronotum brown, thickly yellow pollinose; a small brown median spot at the caudal margin of the scutum. Mesonotum, praescutum dull yellow sericeous, a broad, light brown median stripe, overlain by a dark brown stripe, whose margins are very irregular; two interrupted brown stripes on either side of the median mark, the outermost very pale on the margin of the sclerite; scutum dull brown, with four brown stripes, continuations of the lateral praescutal vittae; the two stripes on each side unite at the caudal margin of the sclerite and run half across the scutellum; scutellum very pale, whitish yellow, sending a median prolongation cephalad onto the scutum; postnotum brown. Pleurae light brown, thickly pale yellowish pollinose. Halteres very pale yellow, the knob brown. Legs: coxae, trochanters and femora dull yellow, the femora darkening to brown apically; extreme base of the tibiae whitish, rest of tibiae and the tarsi dark brown. Wings, veins brown, except costa, which is light yellow and black alternated; membrane hyaline, costal cell with