

Warbler. Rather common during migrations and may breed as a male in full song was shot June 12th, 1882.

38. *Dendroica pinus*; Pine-creeping Warbler. Very rare. Two specimens taken.

39. *Dendroica palmarum hypochrysea*; Yellow Red-poll Warbler. Rather rare during migrations; several specimens taken.

40. *Siurus auricapillus*; Golden-crowned Thrush. Common. Breeds.

41. *Siurus nerius*; Water Thrush. Less common than the last but breeds regularly.

42. *Geothlypis Philadelphia*; Mourning Warbler. Not uncommon in low timbered bush and swampy thickets where it breeds. Marked very rare by McIlwraith.*

43. *Geothlypis trichas*; Maryland Yellow-throat. Common. Breeds.

44. *Myiodytes mitratus*; Hooded Warbler. Very rare. Only one specimen taken.

45. *Myiodytes Canadensis*; Canada Flycatcher. Rather common. Breeds regularly.

46. *Setophaga ruticilla*; Redstart. Common. This species and *D. ustula* are our most common Warblers; this in the wilder and more wooded portions, while the other comes into the towns and inhabits scrub growth in the country.

47. *Vireosylva olivacea*; Red-eyed Vireo. Very common. This species and *V. gilva* are our most common Vireos.

48. *Vireosylva gilva*; Warbling Vireo. Very common. Breeds.

49. *Lanius flavifrons*; Yellow-throated Vireo. Common. Breeds.

50. *Lanius solidarius*; Blue-headed Vireo. Very rare; only one taken September 15th, 1877.

51. *Lanius borealis*; Great Northern Shrike. Rather common in spring and fall. Remains in mild winters but very few breed.

52. *Lanius ludovicianus excubitorides*; White-rumped Shrike. Rather common. Breeds.

53. *Ampelis garrulus*; Bohemian Waxwing. Appears in winter; sometimes in large flocks.

54. *Ampelis cedrorum*; Cherry Bird. Common. Breeds. Remains through mild winters.

55. *Progne subis*; Purple Martin. Common in towns, seldom seen in the country.

56. *Petrochelidon lunifrons*; House Swallows. Breeds abundantly.

57. *Hirundo erythrogastra*; Barn Swallow. Even more abundant than the last.

58. *Tachycineta bicolor*; White-bellied Swallow. Breeds commonly both in town and country.

59. *Cotyle riparia*; Bank Swallow. Breeds abundantly.

60. *Stelgidopteryx serripennis*; Rough-winged Swallow. Breeds in the same localities as the last, but is much less common and generally alone or in small communities by itself. Not given by McIlwraith; probably overlooked.

61. *Pyrranga rubra*; Scarlet Tanager. Common. Breeds. Frequenting mostly pine and other deep woods. Near the shores of Lake St. Clair, it feeds much on the ground.

62. *Hesperiphona ceryleina*; Evening Grosbeak. At Winnipeg, Man., Prof. Macoun was shown specimens of this bird shot near London. McIlwraith gives several shot at Woodstock in 1863.

63. *Pinicola enucleator*; Pine Grosbeak. Occurs quite regularly in winter.

64. *Carpodacus purpureus*; Purple Finch. Common in fall, winter and spring. Breeds sparingly.

65. *Loxia curvirostris Americana*; Red Crossbill. Occasionally common in winter, One shot July 5th, 1882.

66. *Loxia leucoptera*; White-winged Crossbill. Occurs in winter but neither so abundantly nor as often as the preceding.

67. *Agelolus linaria*; Redpoll. Common in winter, arriving in October and leaving in April. A nest of this species with two added eggs was found May 29th, 1879, and identified by comparison with European eggs.

68. *Anthus tristes*; Yellow Bird. Very common, remaining with us in mild winters. Breeds in the third week in July.

69. *Chrysomitris pinus*; Pine Finch. Rather common, arriving in October and leaving in April.

70. *Plectrophenax nivalis*; Snowbird. Common in large flocks in winter; in spring they sometimes stay till late in April and are then in song.

71. *Centropus lapponicus*; Lapland Longspur. First found this spring (1882), at Mitchell's Bay, Kent County, in flocks of from 25 to 200. According to the inhabitants these birds breed there but none were observed after May 18th.

72. *Passerculus sandwichensis nevadensis*; Savannah Sparrow. Regular but not abundant breeder.

73. *Poorella graminea*; Grass Finch. Very common. Breeds.

74. *Coturniculus passerinus*; Yellow-winged Sparrow. Very rare. One specimen taken.

75. *Chondestes grammacus*; Lark Finch. Breeds but is rare. Has been seen common along the roadsides in the southern and western counties.

76. *Zonotrichia leucophrys*; White-crowned Sparrow. Common for a few days in spring, not yet observed in fall.

77. *Zonotrichia albicollis*; White-throated Spar-

* Mr. McIlwraith, of Hamilton, O., published his list of Birds in the "Proceedings of the Essex Institute, Salem, Mass., 1886."