184 MACKAY-THREE CASES OF DOUBLE SALPINGO-OOPHORECTOMY.

Day report, Nov. 9th.—Abdomen greatly distended with flatus and patient very restless. She had a wild and anxious look with eyes somewhat sunken. She looked very bad. Her pulse was 160 and respirations 46, and temperature 100.4. Ordered high enema of soap suds with turpentine half ounce: bowels moved slightly; feeling sick at stomach. Repeated the enema in two hours; bowels moved a little. Patient still greatly dist-essed. At 5 p. m. inserted the rectal tube with good effect. Patient somewhat relieved. Rested very little through the day. Pulse, temperature and respirations unchanged.

Night report.—Eight o'clock, resting nicely. Slept only an hour and a quarter; was troubled very much with tympanites and nausea. Used rectal tube once, with fairly good effect; was given two grains of calomel at 11 p. m. and at 1 and 4 a. m.

Day report, Nov. 10th.—The calomel had good effect; rested well through the day; slept 5 hours. Pulse, 158; temperature, 100; respirations, 38. Ordered peptonized milk by the mouth for first time.

Night report.—Patient had a very good night; slept fully five hours. Felt quite comfortable and refreshed this morning. Patient looked very much better.

Day report, Nov. 11th.—Patient still keeping better; removed part of gauze packing; had slight attack of diarrhæa. Felt quite comfortable. Pulse, 138, and respirations, 32; temperature unchanged.

Night report.—Patient slept very well all night and felt very much better in the morning; had diarrhœa; washed rectum out with weak boracic acid lotion.

Nov. 12th.—Condition the same as on previous day. Diarrhœa better; slept all night; pulse down to 120.

Nov. 13th.—All the gauze packing in pelvic cavity removed. It was perfectly sweet; no discharge; washed vagina thoroughly with bichloride solution, 1 in 5,000. Slept all night; had slight diarrhœa through night.

Nov. 14th.—Patient feeling much better this morning. At 9 p. m. had a slight chill; pulse 160, temperature 102, respirations 38; had pain in lower corner of wound which looked very angry and inflamed. On removing the three lowest stitches quantities of milky fluid welled up from pelvic cavity which was perfectly odorless. A similar discharge came from vagina. Slept fairly well till 4 a. m., but was restless in her sleep.