THE TREATMENT OF CARBUNCLES.

BY THOMAS PAGE GRANT, M.D., LOUISVILLE, KY.

Taking a knife I made a free incision across the top of the carbuncle; after evacuating as well as I could, I washed it out with a solution of carbolic acid about three to five per cent. After this with a pair of dressing forceps I removed all the broken-down tissue I could, a plan which I have found to be of great service in many cases of carbuncles, as thereby whole colonies of micro-organisms are taken out that otherwise would increase and multiply until thrown off by suppuration. Having cleansed the wound thoroughly, I packed it with dry protonuclein special; after which I applied a poultice of flaxseed meal, on which was a teaspoonful of fluid extract of eucalyptus globulus.

As a tonic I ordered:

R

Elix. Ferri, Quiniæ, et Strych. Phosphat., 3 jv

Sig. A teaspoonful three times a day.

The local treatment was repeated for two days, when the poultice was left off and instead this ointment was used:

\mathbf{R}
Sebi ovis)
Sebi ovis) aa
Ceræ flava,
Zinc. oxid.,
Ext. eucalypt. glob
Acid carbolicgrs. c

M. Fiat. unguent. Sig. Grant's Comp. Zinc Ointment.

I continued to wash the wound with the dilute carbolic acid and pack it with the protonuclin; this dressing was renewed twice daily. So rapid was the recovery that on the following Monday evening the wound was healed, and the induration was almost entirely gone, and I dismissed the case with directions that he keep a dressing of the ointment on the seat of the carbuncle for several days to protect the tender skin.