

STACHYS, L. Woundwort.

1800. *S. PALUSTRIS*, L.

Wet ground along streams. July—1. (B.) A variable plant in which 1801 should probably be included as a variety: Plants which answer to var. *cordata*, Gray, and var. *glabrata*, Gray, are found both at Billings Bridge and Kettle Island.

1801. *S. ASPERA*, Mx.

S. palustris, L., var. *aspera*, Gray.

With the last. A slenderer plant with narrower leaves, the angles of the stem beset with stiff reflexed bristles.

PLANTAGINACEÆ.—Plantain Family.

PLANTAGO, L. Plantain.

1804. *P. major*, L. (Common Plantain.)

Introduced. July—1. (B.) The var. *bracteata*, Macoun, is an accidental form which is sometimes found in low ground, but which is not permanent under cultivation.

1805. *P. RUGELII*, Decaisne. (Pale Plantain.)

Low ground. Common. June—4. Easily distinguishable from *P. major* by its slenderer spikes, and much more erect pale yellowish-green leaves. Petioles purplish at the base.

1810. *P. lanceolata*, L. (Rib-wort Plantain.)

Introduced with seed of lawn grasses. Not very persistent. August—2.

1814. *P. media*, L. (Scented Plantain.)

Sparingly introduced; but thoroughly naturalised and very persistent on Parliament Hill and Major's Hill. Leaves ovate, canescently downy, flat on the ground, petioles very short. Flowers silvery and showy with pale purple filaments. Sweetly scented.