

and attributed her suffering in body and mind to this fact. She was despondent, and on the verge of committing suicide. The reflex efforts of the uterine disturbance were also manifested by the derangement of function in nearly all the organs. There was entire loss of appetite and a practical cessation of digestion, accompanied by pain after eating. In October, 1901, I began to give her two capsules of Ergo-Apiol (Smith) three times a day until after her expected periods, without any effect. During the month of November I gave her two capsules three times a day, and continued the treatment until December 12th, 1901, when her menstruation returned in a perfectly normal manner. No unpleasant after-effects whatever were noted at any time during this treatment. She improved both mentally and physically during the time of taking this emmenagogue, and her condition was so remarkably ameliorated that she was discharged cured when the menstrual function had been re-established.

CASE II.—Miss M. B. S. Aged twenty-four years. Has been suffering from amenorrhea for a year, which persisted in spite of all treatment. She was melancholy, and had a very poor appetite and other disturbances due to her suppressed menstruation. In November, 1901, I began giving her two capsules of Ergo-Apiol (Smith) three times a day. I continued this treatment without any appreciable effect, except that the patient seemed to feel more comfortable, and at times during the month she experienced the subjective sensations accompanying the onset of menstruation. Finally, her menses returned on April 21st, 1902. The menstruation was perfectly normal. One week before the next succeeding period I gave her two capsules of Ergo-Apiol (Smith) three times a day, and when the time came for the onset of the flow it appeared in a normal manner. The remedy was continued in doses of one capsule three times a day while the flow lasted. Since the re-establishment of her normal function the patient has gained both mentally and physically, and regained her mental balance and her usual cheerfulness, so that she was discharged cured.

CASE III.—Miss L. D. C. Aged fifteen years. A girl of fine physique, who had first menstruated at the age of nine years, but always very irregularly. The menstruation disappeared for a year and then returned. When admitted she was very irregular with a scanty flow that lasted but one day, and was accompanied by severe pain in the head, loins and pelvis. A week before her expected period in January, 1902, I began giving her one capsule of Ergo-Apiol (Smith) three times a day. At the end of one week her menstruation returned, and lasted four days, the flow being normal in amount and accompanied by very little pain. The same treatment was