Loven for the better Guidance of Shipe, falling in and out of Halfas Halfas Mater. Colour of the Bueys.

Mere placed,

Litchfield Rock

Mars Rock

Detween Chedabucto and si — Ditto

the Litchfield Rock,

Point Pleafant Shoal, the

S. E. part of it,

Shoal between Cornwal-

lis and George's Island

The following is extracted from the GENTLEMAN GARBNER'S DIRECTOR, for the Benefit of those, who are deficus to graft and lay in FRUIT TREES, &c.

Red.

In

ip ii

will

ter t

mes;

Fru

bre

ĊO

To NGUE-GRAFTING, or a Way of grafting in Roots, is thus: The Root is cut stoping about an Inch in Length, and the Graft as much, both very smooth; then each being cleft an Inch is Length also, they are made to enter into one another, so that the Sap of the Graft may join to that of the Root as much as can be. Lap the jointed Part about with a little Hemp or Flax Hurds, and set the Roots so grafted into the Ground, about to or 12 Inches, so, at least, that the Joint may be 4 Inches covered under the Rarth, and may not be bared at any Time, but kept moist by the Rarth. The Root you graft on may be of an Apple or Pear-Tree, about 6 Inches long; it should not be less than the Graft, but it may be bigger, but it is bost when they are both of a Size.

Leying of Trees. This Operation is performed by taking a Bough and laying it into the Ground about fix Inches deep, in good fresh Mould, letting the Top stand about 14 or 16 Inches above the At M. Ground, and keeping the Layers moist during dry Weather: In 8 or 10 Months Time they will take Root. Others cut a Shit in the Layer as in Cornations, then peg down: Others twist the Layer, and Place that is to be covered, like a Withy; and peg down At Hall earth. Everyreses are laid in August; other Trees in

Amn Apolics Inder; if of a small Kind, sive or fix.

er bloffom first at the End of the preceding Year's I if those are shortened, the Bloffom Buds will be taken