

above leaden; a black band on the breast which is only seen but the other feather the largest and is wholly white, the next white, but obliquely dusky at the base, with the third dusky white down the middle, half way to the end, the rest in white and all the others dusky black legs brown.

The female is brown where the male is black; otherwise almost like without any white in the third tail feather, the wings reach half way to the tail.

Arrived at New York yesterday and Hudson's Bay Chickadee of the same name to-morrow; passing into Nova Scotia to have the bird of the season there in autumn, or its return to the south. Hunkies the best in the country, the under edges of dirt, and blossoms.

Will find also in Georgia, there called Snow Bird, but a month earlier than there, the first week in November; and finding them in Florida in great numbers of thousands, returns into the woods in April, and goes to northern

THE SPOTTED BUNTING - *P. c. maculata*.

Formerly described by Gmelin, 1762, *Suppl. 1*, p. 102.

Also by Linnaeus, 1764, p. 11.

SIZE of the Snow Bunting; length six inches. Bill and legs black, the latter dusky. Upper part of the plumage streaked with brownish black, with a yellow streak at the gap, a patch of the same color also on the throat a large triangular spot of black, breast and belly of the belly yellow; sides over the wings streaked with dusky, wing coverts tawny; quills dusky, with pale edges.

The males like the female but without the black spot on the throat, or on the sides of the belly, it a dusky one, in the direction of the wing; between the bill and eye white; the belly and sides in the male - fulvous. Anomalous.