

WHITE-BELLIED SWALLOW. A large wave appeared at Stephenville on June 3rd. The next day they disappeared altogether.

BLACK AND WHITE WARBLER. A few specimens seen in damp evergreen woods.

YELLOW WARBLER. Fairly common. A few nests were located the latter half of June.

MYRTLE WARBLER. Only two birds noted.

MAGNOLIA WARBLER. On June 8th, a nest was located two feet up in a small spruce. The bird had not started to lay. Fairly common.

BLACK-POLL WARBLER. This bird's weak song was heard all along the Humber River in June.

BLACK-THROATED-GREEN WARBLER. One nest found on June 13th contained four fresh eggs. It was placed eight feet up in a pine tree.

OVEN-BIRD. Heard singing in many localities in mixed woods.

WATER THRUSH. Abundant. A nest was found under a bank along the Humber River. On June 12th it contained four fresh eggs.

MARYLAND YELLOW-THROAT. Common.

CANADIAN WARBLER. A moderate summer resident.

AMERICAN REDSTART. Saw several males near Bay of Islands.

AMERICAN PIPIT. On June 19th, a nest with three fresh eggs was found on the side of a moss-covered rock. The nest was built of dead grasses only.

WINTER WREN. Heard everywhere in spruce woods.

RED-BREASTED NUTHATCH. Common.

RUBY-CROWNED KINGLET. Abundant. During the second week in June several nests were found in small spruce trees, eight or nine eggs being the complement in each case.

VEERY. A few individuals noted.

OLIVE-BACKED THRUSH. Very common. Many nests found middle of June on stumps and in spruce trees, three and four eggs forming the set.

HERMIT THRUSH. Heard everywhere during the daytime and night.

ROBIN. Common.

DEPLETION OF BIRD LIFE IN NEWFOUNDLAND.

While the water-fowl are being needlessly slaughtered and exterminated on the sea coast, the birds in the interior are meeting with the same fate. Near two section-houses up on the "barrens" Ptarmigan feathers and skins were scattered about along the railway for some 200 yards. A few casual remarks elicited the fact that one party had shot over 100