

507. *Icterus galbula*, Baltimore Oriole. S. c. B.
509. *Scolecophagus carolinus*, Rusty Blackbird. M. a. It probably breeds sparingly.
- 511b. *Quiscalus quiscula ceneus*, Bronzed Grackle. S. a. B.
- FRINGILLIDÆ—FINCHES, SPARROWS, &C.
515. *Pinicola enucleator*, Pine Grosbeak. W. Irregularly abundant. It appeared in immense numbers in the winter of 1882-3, and again in 1888-9, as did many other of our winter birds.
517. *Carpodacus purpureus*, Purple Finch. S. c. B. Abundant in migration. There are a few winter records of this species, one of which is 29th December, 1885.
521. *Loxia curvirostra minor*, American Crossbill. W. c. Summer records are as follows:—10th May, 1882; 4th August, 1887; 19th June, 1889; 3rd July, 1890.
522. *Loxia leucoptera*, White-winged Crossbill. W. c. A large flock was seen near Beechwood Cemetery in June, 1882.
- 527a. *Acanthis hornemanni exilis*, Hoary Redpoll. W. r. Specimens of this bird, taken by Mr. W. L. Scott in the spring of 1883 were identified by Dr. Coues. It is also included in the list of arrivals for 1887 on March 19th.
528. *Acanthis linaria*, Redpoll. W. a. Summer records are 6th June, 1882; 3rd June, 1888; 22nd May, 1890.
529. *Spinus tristis*, American Goldfinch. S. a. B. It occasionally winters here in large flocks. It did so in 1888-9.
533. *Spinus pinus*, Pine Siskin. W. c. Somewhat irregular in its visits like most of our winter birds. Summer records are as follows:—10th May, 1882; 15th May and 15th August, 1884; 2nd May, 1888; 16th May, 1890.
- *Passer domesticus*, European House Sparrow. R. a. B. Introduced. An intolerable nuisance.
534. *Plectrophenax nivalis*, Snowflake. W. a.
536. *Calcarius lapponicus*, Lapland Longspur. M. c. This species was first recorded here in the spring of 1890, when, in company with Horned Larks, (*O. alpestris*), and Snowflakes, it remained in flocks till May 25th. It was again present in the fall from October 3rd to November 18th.