

weight is increasing. *Lungs*—no dulness or rales, only slight roughening of the breathing.

*Case X.*—J. H., age 22, came under treatment August 14th 1904. He had been coughing and expectorating for seven years. At time of examination cough was troublesome, and expectoration was profuse and purulent. There was marked emaciation, and a chronic purulent notitis media.

*Lungs.*—Softening in the left apex. Patient was put on iodine gr. one half three times a day and cod-liver oil and sent home to the country. At home he had an attack of pleurisy, which seemed to get worse steadily, so he returned to the city.

*Lungs.*—Marked softening in the left apex. Fluid in the left base, and an asthmatic bronchitis. Heart enlarged. Pulse 84, temperature 98. Finger nails were blue. He was emaciated, and suffered much from dyspnoea. Patient was injected with 24 grains, and sent home. His cough improved and he slowly gained in weight. Since that time he has only received 8 grains, as after twenty-four grains it took too long to pick up, owing to the fall in blood pressure and a weak heart. The disease has slowly yielded. The last injection was given January 3rd 1906. After it the patient picked up and had a good appetite. The condition in the lung improved. Dulness has disappeared from the apex, and there were only a few rales in the base. He was put on iodine gr. one half, every four hours and Easton's syrup and went back to the country. Latest report is that his ear has stopped running, and that he was able to eat well, and had only very little cough.

*Case XI.*—Mrs. C., age 24, came to the Montreal Dispensary September, 23rd, 1904. She complained of emaciation and cough, followed by purulent expectoration, and hæmorrhage from the lungs. The weight was 144 lbs. About six months ago she began to cough, and expectorate. Two months ago she had a hæmorrhage from the lungs, which lasted for one week. Since that time cough and expectoration have increased. Sputum contained tubercle bacilli. Patient had two sisters who died of consumption, and one has been treated by this method.

*Lungs*—softening of the left apex. She was put on iodine gr. one-half every four hours and Easton's syrup. Cough stopped October 11th, 1905. Examined January 9th, 1906, in good health and lungs are perfectly normal. Weight 144 lbs. There was no injection in this case, and the results I attribute in part to the iodine.

*Case XII.*—Mr. M., age 20, came under treatment January 1st, 1906. He complained of cough, followed by expectoration, emaciation