

the extreme ultra-violet of certain metals, SEH and D. S. FEHLER	0.25
ion of helium by charcoal, by STUART	0.25
sorption spectra of thallium, aluminium, lead and arsenic, by J. C. McLENNAN, J. F. T. H. J. C. BRETON	0.25
cal transparency of certain specimens of F. T. YOUNG and H. J. C. BRETON	0.25
the Jamin Interferometer for the estimation units of helium or hydrogen in air, by J. C. and R. E. LEWORTHY	0.25
tion of the helium content of mixtures of use of a Katharometer, by V. F. MURRAY	0.25
stability of balloon fabrics to hydrogen and to R. E. LEWORTHY and V. F. MURRAY	0.25
is flow apparatus for the purification of um mixtures, by E. EDWARDS and R. E.	0.25
tion of helium from the natural gases of J. C. McLENNAN	0.25
n of the vapour and liquid phases of the thane-nitrogen, by H. A. McTAGGART and	0.25
line ultra-violet spectra of magnesium and y J. C. McLENNAN, J. F. T. YOUNG, and BRETON	0.25
of light by thin films of rubber, by E. R. T.	0.25
tion of gases by carbonized lignites, by McLENNAN	0.25
y of adsorbing materials, by STUART	0.25
permeability of thin fabrics and films to and helium, by J. C. McLENNAN and W. W.	0.25