

103. **Red-tailed Hawk.** *Buteo borealis*. Common summer resident of the wooded regions. Apparently complementary of the Swainson Hawk, which is found in more open country and on the prairie. Arrives April 15; departs October 15.
- 103a. **Krider Hawk.** *Buteo borealis krideri*. Three specimens of this beautiful Hawk have been taken at Winnipeg--one shot at Rosenfeld by Charles Stewart, Sept. 20, 1905; two now in the collection of A. Calder.
104. **Red-shouldered Hawk.** *Buteo lineatus*. R. H. Hunter writes me that he found this species in Eastern Manitoba.
105. **Swainson Hawk. Common Henhawk** *Buteo swainsoni*. Very abundant summer resident of the prairie region; breeds perhaps twice each season. Have seen several black specimens. Arrives April 15; departs October 15.
106. **Broad-winged Hawk.** *Buteo platypterus*. Regular summer visitant wherever there is woods. In my collection are two specimens, one collected near Winnipeg May 3, 1905, by Ashley Hine. It is reported from various parts of the Province where well timbered, and generally distributed, though not abundant. A. Calder has a beautiful melanistic specimen, killed at Winnipeg, April, 1907.
107. **American Rough-legged Hawk.** *Archibuteo lagopus sancti-johannis*. Migrant; only seen in spring and fall, but not numerous.
108. **Red Roughleg, Gopher-hawk.** *Archibuteo ferrugineus*. One in the collection of E. W. Darbey was shot, in 1895, at Niverville, fifteen miles south-west of Winnipeg, by Geo. Grieve. There is another specimen in the Manitoba Museum.