TREATMENT: In young animals, when cutting teeth, let nature take its course, but when an animal is five years or over, place two drams of Aloin, and two drams of Pulv. Ginger, in a gelatin capsule and administer with capsule gun. Then tone up the digestive organs by mixing one ounce of Pulv. Gentian Root, one ounce of Pulv. Nux Vomica, four ounces of Bicarbonate of Soda. Make into eight powders and give one powder in feed twice daily, or place in gelatin capsule and administer with capsule gun.

LEECHES

(Hæmopis)

The leeches which suck the blood of the horse may be divided into two classes, the external parasites which attach themselves to the skin of the legs and adjacent parts of the horse, and the Hæmopis Sanguisuga, and others of this class, which, not being able to penetrate the skin, endeavor to enter the mouth or nostrils of the horse when he is drinking or grazing in wet and leech-infected pastures. They sometimes cling to the mucous membrane of the eyes. The horse leech, which lives in the water, usually gains access to the mouth and nostrils of the animal, when young and not more than one-tenth of an inch long. They rarely go beyond the air and food passages, generally fastening themselves to the walls of the windpipe and gullet, where they cling till the animal dies from loss of blood or suffocation. They often cause bleeding from the mouth and nostrils, and may be seen by close examination.

TREATMENT: Endeavor to build up the condition of the animal with suitable food. Also feed liberal quantities of stock salt. Where the leeches cling tightly to the mucous membranes of the mouth and nostrils, it is well to cause the horse to inhale the vapor from hor water containing turpentine.