

of the wood. The coarser splinters are *not* detached entirely from the wood, but are split up by the larva all around the top of the cell, and project like bristles, appearing somewhat as those wooden toy-trees which are made for children, and which are formed by shaving down the wood and leaving the shavings still adhering by one end. These splinters make a firm wad. Against them are piled a quantity of finer chips or thin filings, which are loose, but pressed together.

The cell is about 40 mil. from the outer bark of the tree, and the chrysalis makes its way to the air through the burrow by means of its teeth on the segments and the spinose process on the front, by which it forces itself, by stretching and contracting the abdomen, through the wood scrapings which close the cell, until it comes to the end. We have noticed a fine thread of silk proceeding from the spinneret of the larva, although in the puparium we have found no silk whatever. The puparium seems to have been formed by wedging first coarser, and then finer strips of the wood together, and seems to be merely a more carefully and smoothly finished enlargement of the original burrow.

A specimen of the pupa which I have examined is about 30 mil. in length, narrow, brownish black, shining, rugose. The clypeus presents a strong broad spinous process, supported at base by lateral projections. On the underside it descends into a wide sulcation, terminating in a broad projection. The capital appendages are visible, and here and there arise isolated hairs as in the previous stage. The abdominal segments are provided with teeth over the dorsum, decreasing in size to the stigmatal line. The anal segment is provided with two unequal sized terminal teeth on each side of the vent.

The moth seems to belong to the genus *Cossus* Fabr., and not to be congeneric with *Xystus robinia*. The head is short, eyes naked, labial palpi small, appressed, scaled. The thorax is thickly scaled, the scales gathered into a ridge behind, and is squarer in front than in *Xystus*, not so elongate, or so elevated dorsally. The male antennæ are bipectinate, the lamellæ short, rather broad and ciliate. The female antennæ are serrated. It is allied to the European *Cossus terrebra*, while a larger insect. It differs from *C. querciperda* by the absence of any yellow on the male hind wings, and by its darker color and closer reticulations.

In color this species is black and gray; the edges of the thorax and collar are shaded with gray—more noticeable on some specimens than others. The primaries are covered with black reticulations, which are not