

fecal ptomaines and leucomaines a prominent factor in originating anæmia and chlorosis occurring in young women of constipated habit. Until recently notices had not been taken of acute fecal poisoning. Burford *Lancet* reports 11 cases of mild septic toxæmia following the use of copious warmed enemata to relieve the constipation of women under treatment for uterine disease. The cases related form an ascending series. The fact displayed a generalized rash and congested throat, but no pyrexia. In the eighth there was a diffused erythematous rash, a severe sore throat, and a temperature of 102°. The duration of the attack varied between two and four days. Liquefaction of the feces is promoted by the enema. During its temporary retention osmosis occurs, lymphatics and blood vessels readily absorbing the warm septic material. Three to four per cent. of the enemata administered produced the condition narrated. Burford found it especially liable to occur if enemata were used within three days subsequent to the administration of ether by inhalation in cases in which free evacuation of the bowels had not occurred previously. The character of the fluid used did not especially influence the production of the septic condition. It is more likely to follow large than small enemata.—*Poly-clinic*.

# REFERENCE TABLE OF DOSES.

BY JOSEPH W. ENGLAND, PH. G.

(Concluded.)

DRUG.	DOSE.
Theina	1-2 3 gr
Theinæ Citras } Max	3-5 gr
Thymol	3-5-10 gr
Tinct. Aconiti	1-2-3 m
Max	3-5 m
Tinct. Aloes	1-2 4 fl dr
" Aloes et Myrrhæ	1-2 fl dr
" Asafetidae	1-1 1/2 fl dr
" Belladonnae	5-10-15 m
Max	15-20 m
" Benzoini	10-15-30 m
" Benzoini Comp.	15-30-60 m
" Calumbæ	1-2 4 fl dr
" Canab. Indic.	10-15-30 m
Max	30-60 m

DRUG.	DOSE.
Tinct. Cantharidis	3-5-10 m
" Capsici	10-30-60 m
" Cardamomi Comp	1-2 3 fl dr
" Catechu Comp.	1-3 1 fl dr
" Chiratae	1-2 fl dr
" Cinicifugæ	1-2 4 fl dr
" Cinchonæ	1-2 3 fl dr
" Cinchonæ Comp	1-2 4 fl dr
" Colchici Rad.	5-15-30 m
Max	30-60 m
" Colchici (sem)	10-30 60 m
Max	1-1 1/2 fl dr
" Conii	15-30-60 m
" Cubebæ	1-2 fl dr
" D gitalis	5-15-30 m
Max	30-60 m
" Ferri Acetatis	10-30-60 m
" Ferri Chloridi	5 10-20 m
" Gallæ	1 2 fl dr
" Gelsemii	5 10 20 m
" Gentianæ Comp	1 2 4 fl dr
" Guaiaci	1 2 fl dr
" Guaiaci Amuon.	1 2 fl dr
" Hellebori	15 30 60 m
" Humuli	1 2 3 fl dr
" Hydrastis	1 1 fl dr
" Hyoscyami	10 30 60 m
Max	1 2 fl dr
" Ignatiæ	5 10 15 m
Max	15 20 m
" Ipecac. et Opii	5 10 15 m
" Jalapæ	1 2 fl dr
" Kino	1 3 fl dr
" Krameriæ	1 3 fl dr
" Lavandulæ Comp	1 2 fl dr
" Lobeliæ Expect	10 30 60 m
Emet. (Asth.)	1 2 fl dr
" Lupulinae	1 2 fl dr
" Matico	1 2 fl dr
" Myrrhæ	15 30 60 m
" Nucis Vomicae	5 10 20 m
Max	20 30 m
" Opii	5 10 15 m
Max	15 30 m
" Opii Acet.	5 10 15 m
Max	15 30 m
" Opii Camphor	1 2 4 fl dr
" " Deodor	5 10 15 m
Max	15 30 m
" Physostigmatis	10-15-30 m
" Quassiæ	1-1 1 fl dr
" Rhei	1-2 4 fl dr
" Rhei Aromatica	1-1 1/2 3 fl dr
" Rhei Dulcis	1-2 4 fl dr