



FIG. 5.

than broad, fifth about one and one-half times as long as broad at tip, with two long hairs at tip, the outer one more than twice as long as joint; body almost twice as long as broad, sides (beyond eyes) sub-parallel, broadly rounded behind; two eyes each side, and near them a long seta; two setae in front and two on middle of the cephalothorax; on anterior margin of the cephalothorax is a single black quadrangular mark with a white circle in it. Length, .75 mm. Washington, D. C. (March.)

Specimens were taken on wet ground under stones and among short grass; it is the first species of the genus that I have seen from America.

*Notophallus dorsalis*, n. sp.—Black; legs red; a large elongate red spot on the posterior median dorsum, containing the anal opening; a large eye each side on the anterior portion; some scattered short hairs above. Legs short, first pair scarcely longer than the body, fourth pair shorter than the body. Genital opening large, elliptical, divided longitudinally; palpi short, red. Length, .8 mm.

A few specimens taken at Washington, D. C., in the early spring; on ground under stones and sticks. It is the first record of the genus in this country; a genus differing from all our other Eupodidae in having the anal opening on the dorsum.

*Cheyletus clavispinus*, n. sp.—Body pale reddish or yellowish, fading out in alcohol. Body somewhat quadrangular, corners rounded, about