

Lepeophtheirus robustus Krøyer.

Lepeophtheirus robustus, KRØYER, H. "Snyltekrebsene." Naturhistorisk Tidsskrift, ser. 3, vol. 2, 1863, p. 135; pl. 6, fig. 6, *a-c*.

Found on the gills of various rays from the Greenland coast.

Lepeophtheirus salmonis Krøyer.

Lepeophtheirus salmonis, WILSON, C. B. "Pacific Coast Copepods." Proc. U. S. Nat. Museum, vol. 35, 1908, p. 419.

From the outer skin of the hump-backed salmon, *Oncorhynchus gorbuscha*, and the Dolly Varden trout, *Salvelinus malma*, at Karluk, Alaska.

Lernæocera branchialis (Linnæus).

Lernæa gadina, FABRICIUS, OTTO. "Fauna Groenlandica", 1780, p. 336; No. 325.

Lernæa branchialis, STEPHENSEN, K. "Grønlands Krebsdyr." Meddel. om Grønland, vol. 22, 1913, p. 318.

Found on the gills of various Gadidae from the Greenland coast.

Lernæocera godfroyi (Quidor).

Lernæa godfroyi, QUIDOR, A. Deuxième Expédition Antaretique Française. "Copepodes parasites." Sci. Nat.; Documents scient., 1912, p. 210; pl. 2, fig. 23.

From the gills of *Cottoperca dollói* (?), at Tuesday bay, Antaretic ocean.

Nesippus borealis Steenstrup & Lütken.

Nogagus borealis, STEENSTRUP, J. & LÜTKEN, CH. "Snyltekrebs og Lernæer." Kongl. Danske Vidensk. Selsk. Skrifter, ser. 5; vol. 5, 1861, p. 387; pl. 11, fig. 21.

Nesippus borealis, WILSON, C. B. "The Pandarina and Cceropinae." Proc. U. S. Nat. Museum, vol. 33, 1907, p. 438; pl. 37.

From the coast of Alaska, host unknown.

Peniculus clavatus (Müller).

Peniculus clavatus, KRØYER, H. "Snyltekrebsene." Naturhistorisk Tidsskrift, ser. 3, vol. 2, 1863, p. 266; pl. 14, fig. 8, *a-g*.

From the fins of *Sebastes norregicus* (Cuv. & Val.), on the Greenland coast.

Pennella antarctica Quidor.

Pennella antarctica, QUIDOR, A. Deuxième Expédition Antaretique Française. "Copepodes Parasites." Sci. Nat.; Documents scient., 1912, p. 206; pl. 1, fig. 15-17; pl. 1, fig. 29-31.

From the skin and blubber of whales belonging to the genera *Balaenoptera* and *Megaptera*, in the antaretic ocean.

Pennella balænopterae Koren & Danielssen.

Pennella balænoptera, QUIDOR, A. Deuxième Expédition Antaretique Française. "Copepodes Parasites." Sci. Nat.; Documents scient., 1912, p. 205; pl. 1, fig. 14; pl. 4, fig. 32.

From the skin of *Balaenoptera sibbaldi*, in the antaretic ocean.

Pennella charcoti Quidor.

Pennella charcoti, QUIDOR, A. Deuxième Expédition Antaretique Française. "Copepodes Parasites." Sci. Nat.; Documents scient., 1912, p. 207; pl. 1, fig. 13; pl. 2, fig. 25; pl. 4, fig. 33.

Found on the skin of a species of *Balaenoptera* in the antaretic ocean.