

in bed, either indoors or in the open air, abundance of milk, massage, and regulated exercises are desirable in the majority of cases.

REFERENCES.

1. Quineke. *Berlin. Klinisch. Wochenschrift.* 1891.
2. Widai. *Rev. Neurolog.* 1903.
3. Nouné. Refd. by Stertz. *Wiën. Med. Klinik*, No. 4. 1912.
4. Mott and Haliburton. *Brit. Med. Journ.* 1904.
5. Mott. *Brit. Med. Journ.* Nov. 8, 1911.
6. Flexner and Lewis. *Journ. Amer. Med. Assoc.* 1909.
7. Turner. "Epilepsy," p. 212. 1907.
8. P. Clark. *Archiv. of Int. Medicine.* 1912.
9. Turner. "Epilepsy." 1907.
10. Freud. *Selected Papers*; trans. by Dr. Brill. New York. 1909.
11. Pawlow. Refd. by Morton Prince. *Journ. of Abnormal Psychology.* 1907.
12. Janet. "Major Symptoms of Hysteria." 1906.
13. Ormerod. *Brain*, 1911.
14. Wilson. *Brain*, 1911, to whose critical digest on *Hysteria I* am indebted for much information.
15. Babinski. *Soc. de Neurol.* 1901.
16. Freud. *Selected Papers.* Dr. Brill's translation. 1909.
17. Dr. Brill's trans., p. 188.
18. Hart. *Brain.* 1911.
19. Mills. *Monthly Cyclopedia.* July, 1908.
20. Dejerine. "Les Psycho-Névroses." 1911.
21. Dubois. "Psychical Treatment of Nervous Disorders." 1909.
22. Freud. *Selected Papers.* Op. cit. Chap. 4.

CARDIAC ASTHMA.

Rosin (*Deutsch. med. Woch.*) says treatment must be rapid and heart tonics should be given by subcutaneous, intramuscular or intravenous injection, as drugs given by the mouth disturb respiration and are defectively absorbed. Camphor in a 10 per cent. oil solution is very useful and indispensable during attacks. The injections can be repeated every fifteen minutes.