

as a result of the global increases in the atmospheric concentrations of greenhouse gases and the associated climatic changes. Projected changes in temperature and precipitation suggest that climatic zones could shift several hundred kilometres towards the poles over the next fifty years. Flora and fauna would lag behind these climatic shifts, surviving in their present location and, therefore, could find themselves in a different climatic regime. These regimes may be more or less hospitable and, therefore, could increase productivity for some species and decrease that of others. Ecosystems are not expected to move as a single unit, but would have a new structure as a consequence of alterations in distribution and abundance of species.

The rate of projected climate changes is the major factor determining the type and degree of climatic impacts on natural terrestrial ecosystems. These rates are likely to be faster than the ability of some species to respond and responses may be sudden or gradual.

Some species could be lost owing to increased stress leading to a reduction of global biological diversity. Increased incidence of disturbances such as pest outbreaks and fire are likely to occur in some areas and these could enhance projected ecosystem changes.

Consequences of CO₂ enrichment and climate change for natural terrestrial ecosystems could be modified by other environmental factors, both natural and man-induced (e.g. by air pollution).

Most at risk are those communities in which the options for adaptability are limited (e.g. montane, alpine, polar, island and

coastal communities, remnant vegetation, and heritage sites and reserves) and those communities where climatic changes add to existing stresses. The socioeconomic consequences of these impacts will be significant, especially for those regions of the globe where societies and related economies are dependent on natural terrestrial ecosystems for their welfare. Changes in the availability of food, fuel, medicine, construction material and income are possible as these ecosystems are changed. Important fibre products could also be affected in some regions.

2.3 Hydrology and water resources

Relatively small climate changes can cause large water resource problems in many areas, especially arid and semi-arid regions and those humid areas where demand or pollution has led to water scarcity. Little is known about regional details of greenhouse-gas-induced hydrometeorological change. It appears that many areas will have increased precipitation, soil moisture and water storage, thus altering patterns of agricultural, ecosystem and other water use. Water availability will decrease in other areas, a most important factor for already marginal situations, such as the Sahelian zone in Africa. This has significant implications for agriculture, for water storage and distribution, and for generation of hydroelectric power. In some limited areas, for example, under an assumed scenario of a 1°C to 2°C temperature increase, coupled with a 10% reduction in precipitation, a 40-70% reduction in annual runoff could occur. Regions such as southern Asia, that are dependent on unregulated river systems, are particularly vulnerable to hydrometeorological change. On the other hand, regions such as the western USSR and western United States that have large regulated water resource systems are less sensitive to the range of hydrometeorological changes in the assumed scenario. In addition