

- BONAPARTE GULL, *Larus philadelphia*—A small flock of ten or fifteen usually to be found during August, feeding in the middle of the lake.
- CASPIAN TERN, *Sterna caspia*—This rare species is one of the notable birds of the lake. Their favorite roost seems to be the stumps half way to Fenelon Falls, but later they were discovered in similar stumpy areas in the marsh. They are also to be found fishing singly or in pairs up and down the lake, and may be distinguished from the gulls even at a distance, by the characteristic, down-pointing head and bill. The identification is made doubly sure when one drops into the water with a great splash to secure some luckless minnow.
- BLACK TERN, *Hydrochelidon nigra surinamensis*—Fairly common. Breeds in the marsh. Early migrant. Nearly all gone by the middle of August.
- BLACK DUCK, *Anas rubripes*—Flocks, sometimes numbering hundreds, seen in the marsh.
- AMERICAN BITTERN, *Botaurus lentiginosus*—Common.
- GREAT BLUE HERON, *Ardea herodias*—Very common.
- GREEN HERON, *Butorides virescens virescens*—Along the river in the marsh. Somewhat rare.
- VIRGINIA RAIL, *Rallus virginianus*—The only one observed was on August 19th, 1914, but no doubt there are numbers of them throughout the marsh.
- SORA RAIL, *Porzana carolina*—Several records for this bird. Also no doubt quite common.
- FLORIDA GALLINULE, *Gallinula galeata*—Very abundant. One of the most interesting birds of the marsh.
- AMERICAN COOT, *Fulica americana*—A few seen each year. Not nearly as common as the Gallinule.
- LEAST SANDPIPER, *Pisobia minutilla*—A small sandpiper observed from the deck of the steamer en route from Lindsay to the Point. It flew up as the boat approached and identification, of course, is not positive.
- SOLITARY SANDPIPER, *Helodromas solitarius*—During the wet summer of 1912 several were noted alongshore and in a pool in a clearing in the woods. None seen in 1913 or 1914.
- SPOTTED SANDPIPER, *Actitis macularius*—Common.
- BLACK-BELLIED PLOVER, *Squatarola squatarola*—August 9th, 1912, one seen in a wet field three miles from the lake—a rather unusual place. Identified by the white rump and black patches under the wings.