

SPRAYING CALENDER.—(Continued).

6. <i>Grape</i>	Copper sulphate when buds swell. Paris green for flea beetle.	Bordeaux when leaves 1½ inches in diameter; Paris green for beetle.	Bordeaux when flowers have fallen; Paris later.	Bordeaux 10-15 days later if disease still appears.*
Mildew, etc., flea beetle.				
7. <i>Raspberry</i>	Copper sulphate before buds break.	Bordeaux if rust appears during summer.	Bordeaux if the trouble appears to continue.	The only remedy, as yet, for orange rust is to cut out diseased plants.
8. <i>Currant</i>	Paris green or hellebore for worms.	Hellebore 10 days later for worms; Bordeaux for mildew.		
9. <i>Gooseberry</i>	Bordeaux as soon as leaves expand for mildew, hellebore for worms.	Bordeaux 10-15 days later. Worms as before.	Ammoniacal copper carbonate 10-15 days later.	If further treatment required for mildew repeat third 10-15 days later.
10. <i>Tomato</i>	Bordeaux as soon as rot or blight appears.	Bordeaux if trouble continues.		
11. <i>Strauberry</i>	Bordeaux when first fruits are setting.	Ammoniacal copper carbonate when first fruits are ripening.	Bordeaux when fruit is taken off.	Bordeaux if trouble continues.
12. <i>Potato</i>	Paris green as soon as beetles appear.	Paris green when plants are about six inches high.	Bordeaux 10-15 days later.	Repeat if necessary in 10-15 days.
13. <i>Cabbage</i>	Pyrethrum may be applied in solution or dusted on, 1 part pyrethrum to 6-8 parts flour.			