

Seasonable directions are given in the *Medical Standard* for the treatment of Summer Diarrhoea: Carharrer recommends the discontinuance of milk as a diet, and substitution of liquid peptonoids with cocoa, pure brandy, and beef tea or broths. Water must be allowed freely, given in small quantities, and often. In bad cases he gives:—

R. Acidi salicylic., gr. ss
Cretæ precip., gr. x
Glycerin, ʒ ij
Aquæ rosæ, ʒ xiv, M.

Sig.—Fluid ounce every hour for a child one year old.

The following mixtures will prove of benefit in the treatment of intractable cases:—

Loomis' diarrhoea mixture.

Tincture of opium, ½ fl. oz.
Tincture of rhubarb, ½ fl. oz.
Compound tincture of catechu (U. S. P.), 1 fl. oz.
Oil of sassafras, 20 mins.
Compound tincture of lavender, enough to make, 4 fl. oz. M.

Sig.—One teaspoonful every four hours, for adults.

Squibb's diarrhoea mixture.

Tincture of opium, 1 fl. oz.
Tincture of capsicum, 1 fl. oz.
Spirits of camphor, 1 fl. oz.
Purified chloroform, 180 mins.
Alcohol, enough to make, 5 fl. ozs. M.

Sig.—One teaspoonful every five hours, for adults.

Thielmann's diarrhoea mixture.

Wine of opium, 1 fl. oz.
Tincture of valerian, 1½ fl. oz.
Ether, ½ fl. oz.
Oil of peppermint, 60 mins.
Fluid extract of ipecac, 15 mihs.
Alcohol, enough to make, 4 fl. ozs. M.

Sig.—Thirty drops every three to five hours, for adults.

Velpeau's diarrhoea mixture.

Take or tincture of opium, compound tincture of catechu (U. S. P.), spirit of camphor, each equal volumes. M.

Prof. Parvin presented to the class a case of *vulvitis* resulting from diabetes mellitus, and directed the following:—

Constitutional treatment for the diabetes and local treatment for the vulvitis, as follows:—

R. Atropin., gr. j
Aquæ, ʒ ʒj. M.

To be used as a spray.

Also an ointment composed of sodium salicylate, benzoated zinc ointment, and tar ointment.

A Paris correspondent of *The Times and Register*, June 7th, gives the modern medicinal treatment for Tonsillitis. As it is now to be considered as an infectious malady, antiseptics are in order. This may be used first, for buccal antiseptics:—

R. Borate or benzoate of soda, ʒ ijss
Hot water, ʒ vij

Dissolve, and add:

Tincture of myrrh, gr. lxxv
Blackberry syrup, ʒ j M.

Ft. gargle.

Or the following:—

R. Resocine, gr. xv
Distilled water, ʒ ʒj
Blackberry syrup, ʒ ʒj. M.

Ft. gargle.

Then brush over the tonsils, several times a day, with the following:

R. Glycerine, ʒ v
Camphor, gr. xv
Carbolic acid, gr. xv M.

Use as above, with camel's-hair brush.—*Coll. and Clin. Record.*

Merck's Bulletin, May, 1890, gives the following formula for a megrin powder:—

Caffeine citrate, true, gr. xv
Phenacetine, gr. xxx
Sugar, gr. xv. M.

Divide into ten wafer-powders. One wafer every two or three hours.

For sweating feet, Legoux (*Nouv. Remèdes*, quoted in *Nat. Druggist*, May, 1890, recommends:—

R. Ferri perchlorid., ʒ ij
Glycerin, ʒ j
Essent. Bergamot, ʒ ij. M.

Sig.—Apply with a pencil or swab.

For Chapped Hands, a writer in the *Provincial Medical Journal* suggests the following as an excellent application:—

R. Menthol, gr. xv
Salol,
Olive oil, āā gr. xxx
Lanolin, ʒ iss. M.

As an external application in acute rheumatism, a writer in the *London Medical Record* suggests the following:—

R. Salol,
Ætheris, āā p. iv
Collodii, p. xxx. M.

In Gonorrhoea, Julien (*Revue de Therap.* March 25th, in *Med. News* April 5th) recommends the following injection:—

R. Liquid vaseline, p. 140
Bismuth Subnitrate, p. 10
Resorcine, p. 3
Iodol, p. 1. M.