

or every third night, by bathing, and changing clothes in about the same way and at about the same interval for a week, a cure is effected in ordinary cases.

The male acarus is not a burrower. At the time the female has proruptured into the external world from below the epiderm, the male is either dead or certainly functionless.

Few cases or case groups last over the ninth day under this treatment.—*Therapeutic Gazette*.

Simple Catarrhal Conjunctivitis.

R̄ Acidi borici	gr. 40.
Sodii chloridi	gr. 6.
Aq. camphoræ,	
Aq. dest	āā ʒ 2.

M. Sig.: Apply as lotion to eye every two hours.—*Jour. Amer. Med. Assn.*

Dyspepsia and Gastro-Enteritis.

R̄ Creosote (beechwood)	gtt. iij.
Alcohol	℥xv.
Powd. acacia	ʒ iiss.
Syr.	ʒ j.
Aq. aurantii flor. fl	ʒ iiss.
Aq., q. s. ad fl	ʒ iij.

M. Sig.: A teaspoonful for children, a tablespoonful for adults, t. i. d., before meals.—*Zangger, Phila. Med. Jour.*

Angeioma.

Blaschko (*Munchener medicinische Wochenschrift, Lyon medical*) recommends the following formula:

R̄ Perchloride of iron.	
Chloride of zinc	āā gr. 15.
Hydrochloride of cocaine	gr. 1½.
Aq. dest	gr. 150.

M. Sig.: From half to a whole Pravaz syringe-ful to be injected every two or three days.—*N. Y. Med. Jour.*

Pain and Tenesmus of Dysenteric Diarrhea.

Taylor and Wells recommend the following analgesic suppositories:

R̄ Cocain. hydrochlorate	gr. j.
Aq. ext. of ergot	gr. xij.
Aristol	gr. v.
Cocoa butter to make twelve suppositories.	

Sig.: One to be applied every hour or longer.—*Columbus Med. Jour.*