

actions upon the human economy are those of foreign bodies, irritating and depressing until thrown off by the conservative and defensive forces of the body.

A VERITABLE PROP.

After the subsidence of the acute symptoms of any serious febrile disease, an examination of the blood will almost always reveal a degree of anemia in direct proportion to the severity and duration of the primary disease. It is thus always desirable in such cases to adopt measures to revive, restore and reconstruct, and with this object in view one should begin at the foundation, i.e., the blood itself. To construct new red cells, and reconstruct those which have become dehemoglobinized by disease, nothing is more potent in effect than Pepto-Mangan (Gude). This standard preparation of organic iron and manganese supplies the vital fluid with the elements needed to reconstruct and restore its oxygen carrying capacity, by contributing the necessary hemoglobin. Pepto-Mangan is palatable, absorbable, and promptly assimilable. It encourages the appetite, without disturbing digestion or causing constipation.

GONORRHOEA IN WOMEN.

The application of an aqueous solution of iodine in connection with the administration of sanmetto is recommended in the treatment of gonorrhoea in women on account of its ability to penetrate the subepithelial structures and deeper glands. It is also a stimulant, a counter-irritant, and an alterative. The aqueous solution (liquor iodi compositus) is used because of the pain caused by alcoholic tincture. The gonococci disappear early from the secretions, intrapelvic extension is far less frequent, and there is little pain. The entire course of treatment is comparatively short and causes but little trouble to either patient or physician. In acute and sub-acute cases, where vaginitis or vulvitis is present, the patient should be directed to use one teaspoonful of the solution to two quarts of warm water as a douche twice daily. In the chronic and mildly inflammatory cases the strength should be gradually raised from one to two teaspoonfuls, or until the patient begins to experience a burning or smarting sensation, indicating the limit of increase, the sanmetto to be administered in teaspoonful doses for times daily throughout the treatment.