

been supplied last saring for seed a large quantity of mustard, thistles, etc., has been imported and a good deal of indignation is expressed in consequence.

The yield reported as likely to be had over the Province is 44.3 bushels per acre. In 1887 the yield as reported by correspondents was 43.1 and in 1889 21.5 bushels. In the North Central group a yield of 47.1 bushels to the acre is expected.

BARLEY.—Like wheat and oats, barley is reported a good healthy crop and is likely to yield handsomely. The growing time has been very favorable and although the spring season was late, still quite a large quantity was harvested during the last few days of July, and the great bulk of the crop would be ready for harvesting during the first week of August. Weeds the result of impure seed, in many places troubled the crop.

DISTRICTS.	APPROXIMATE AVERAGE YIELD PER ACRE.				
	Wheat.	Oats.	Barley.	Prairie Hay.	Grasses, etc.
	Bus.	Bus.	Bus.	Tons.	Tons.
Northwestern	26.9	45.1	34.5	1.77	2.56
Central	24.5	44.5	35.8	1.28	2.01
Southwestern	23.9	44.6	34.7	1.48	2.04
South Central	21.4	42.2	32.9	1.09	1.66
North Central	26.2	47.1	34.5	1.53	1.47
Eastern	24.6	42.3	33.2	1.38	1.41