

FAMILY ICHNEUMONIDÆ.

358. *ICHNEUMON FERRUGATOR Kirby*.—Length of body 7 lines. Taken in the Expedition, but no locality stated.

[259.] Abdomen black, rather glossy, very thickly punctured with minute and often confluent punctures. Head transverse, triangular, not quite so wide as the middle of the trunk; anterior margin of the face rounded; palpi reddish; eyes long, subelliptical; antennæ shorter than the trunk, spirally convoluted; trunk oblong, subcompressed; scutellum subtriangular, rounded at the apex; metathorax armed on each side with a short tooth, with several elevated longitudinal and oblique lines; legs with decumbent whitish hairs, anterior tibiæ obscurely, and all the tarsi, rufous; wings embrowned with a rufous tint, nervures darker; abdomen linear-lanceolate, rufo-ferruginous, with the first joint, which is dilated at the apex, black; footstalk channelled longitudinally on each side.

FAMILY CRYPTIDÆ.

359. *CRYPTIUS VIDUATORIUS Fabr.*—Length of body, with ovipositor, $5\frac{1}{4}$ lines; do, without ovipositor, $3\frac{1}{2}$ lines. One specimen taken in Lat. 54° .

Body black. Head subtriangular, transverse, very minutely and thickly punctured; palpi pale rufous; face plane with two elevations in the middle; eyes large, oval; antennæ shorter than the trunk, rather slender, involute, black with a white band in the middle; orbit of the eyes behind with a very indistinct white line; trunk oblong, cubical, gibbous, very thickly and confluent punctured except on the back; tegulæ white; metathorax armed with two minute teeth on each side, one in the middle and the other at the base, forming the terminal angle, marked out into three areas by elevated lines; legs red with coxæ, trochanters, and posterior tarsi black; wings subtestaceous with the larger nervures black; abdomen, excluding the ovipositor, scarcely so long as the trunk; first segment impunctured, glossy, dilated at the apex, which is subquad-rangular; footstalk flat; second and third segments very large, with their gloss obscured by infinitely minute punctures; ovipositor shorter than the abdomen; borer red.