

recovered from the operations, she declared that she was feeling better than she had been for years. But very soon the fetid discharge and the pain returned at each menstrual period, and evidently something else had to be done if I wanted to save my reputation. I then tried local applications, alteratives, uterine tonics, etc., all without avail, until finally Ergoapiol (Smith) was given. The result was immediate relief and a gradual and permanent improvement in the menstrual flow until it was free from pain and devoid of any disagreeable odor.

This patient was evidently suffering from congestive dysmenorrhea which was intensified by the presence of lacerations of the cervix and the perineum which had existed since parturition. The result attained illustrates very well how Ergoapiol (Smith) acted upon the uterus, restoring its tissues to normal condition and re-establishing the menstrual function upon its normal basis.

Another type of dysmenorrhea, that which I term "nervous," but which the authorities term "neuralgic," is illustrated by the following case which recently came under my care:

The patient was a young woman who had been married two years, but had not borne any children. She stated that she had pain during the menstrual period from the first onset of menses, and at the time of examination she also complained of a fetid discharge. The menstrual flow was scanty and rarely of blood red color. Just before and after the period, she had backache and headache, her complexion was unhealthy, not bright and clear as that of her sister, and she appeared older than she really was. She always dreaded the onset of the menses which recurred with a regularity that is not always present in these cases. She was easily excited, and received impressions vividly and indelibly. Her digestion was poor, and she was often sleepless, irritable, and peevish.

Vaginal examination revealed a slightly thickened os and slight endocervicitis with erosions of the cervix. Cod-liver oil, malt extract, hypophosphites, and aromatics, in combination, 25 per cent. of each, were given freely during the intervals between the menstrual periods, and for three days before the expected menstruation Ergoapiol (Smith) was given in capsules, one being given three times daily until the discharge ceased. At the fourth period after the beginning of the treatment she was relieved of all her symptoms, and was free from pain and fetor during menstruation. Locally, tincture of iodine and occasionally tampons of ichthyol and glycerine were applied. The cure was permanent and in every way satisfactory.