

Publishers' Department

I HAVE used Resinol Ointment as a dressing for slight wounds, minor surgical purposes and for piles. In every case I have found your Ointment to be all it is represented to be—a great curative. I have known instant relief result from its application for piles. I heartily recommend it to my associates as a useful article, and one that is handy to have in many emergencies.—E. STREET, Nurse, "The Sanatorium, Matlands, Australia.

SOME USEFUL SUGGESTIONS.—In urinary incontinence, especially in children, give atropine, gr. 1-500 t. i. d. and an extra granule at bed time. In urinary incontinence due to general weakness, especially of the sphincters, add strychnine nitrate, gr. 1-134 to each dose. In urinary incontinence due to irritation of bladder from foul urine add lithium benzoate, gr. 1 to each dose. In urinary incontinence from cystitis, add a granule each of aconitine (gr. 1-134) and cannabin to each dose. Atropine, gr. 1-500 to 1-250 every fifteen minutes is useful in hemorrhage from any part of body. Hydrastinine, gr. 1-12, should be added to atropine if the bleeding is from uterine membranes. This combination effective. Atropine, gr. 1-250, till dryness of mouth is noticed, is efficacious in ovarian neuralgia, dysmenorrhea, etc. Dioscorein, gr. 1-6, should be added to relieve the severer pains. In excessive sweating, especially at night, pulse slow and soft, skin cold and clammy, atropine, gr. 1-250, every half hour will soon change conditions. In acute coryza, beginning of congestion, give atropine, gr. 1-500 every half hour until dryness of the throat is noticed, then less often. Hamamelin, gr. 1-6, should be added to atropine if bleeding from veins is due to a general relaxation of their walls. I gave dioscorein, gr. 1-6, in hot water every ten minutes, in severe gall-stone colic, and sodium succinate ever since for last six