

provement.—Eleven a.m. : Ordered to discontinue the use of the Calabar bean. To produce continued sleep, half a grain of the acetate of morphia was injected beneath the skin ; also the same quantity at one p.m. and four p.m. respectively. At one p.m. an enema of beef tea and brandy was administered.—Eight p.m. : In a deep sleep.—Ten p.m. : Pupils much contracted ; still asleep.—Forty minutes past ten : He awoke, and drank twelve ounces of strong beef tea and two ounces of brandy. He says he feels better. Pulse 136. Injection of half a grain of acetate of morphia.

7th.—He awoke and took some beef tea and brandy. At ten minutes past two a.m., and again and half past eight and eleven, one grain of the morphia was injected hypodermically.—Forty-five minutes past five p.m. : Has slept continuously. Motions passed of a light color ; has taken nourishment.—Twenty minutes past seven p.m. : Feels more comfortable. The morphia injection (half a grain) was repeated.

8th.—Ten a.m. : Injection of a grain of acetate of morphia ; pulse 128.—Five minutes past two p.m. : The spasms are much diminished ; he lies in a comfortable sleep ; pulse 120, and feeble. Ordered five grains of disulphate of quinine to be administered three times a day at proper intervals.—Quarter past eleven p.m. : Injection of half a grain of the morphia.

9th.—One grain of the acetate of morphia injected. The “snapping” of the jaws diminished, and he moves his arms and speaks with ease.—Five p.m. ; One grain and a half of the acetate injected.—Eight p.m. : One grain of the morphia injected.

10th.—Passed a good night. At fifty minutes past two, one grain, and at a quarter past eight two grains, of the acetate were injected.

11th.—He seems quite comfortable.—Quarter past eleven : Injection (two grains) repeated.

12th.—Convalescent.

22nd.—The medicines have been gradually discontinued.—*Lancet*.

## REPORT OF A CASE WHERE A PORTION OF TOBACCO-PIPE WAS BROKEN INTO THE BLADDER.

### SUCCESSFUL REMOVAL AFTER THE USE OF THE LITHOTRITE.

By HENRY SMITH, F.R.C.S., Assistant-surgeon to King's College Hospital.

J. H—, aged nineteen, was sent to my house on Saturday, Jan. 16th, by Dr. Riding, with the following story : Three months previous, having a difficulty in micturition, he conceived the idea of passing a long clay pipe down his urethra into his bladder. He was so success-