

from attempting the operation, stating that "it was sure to be a failure." Not only did he do this, but used his endeavors to prevent the lady from having the operation performed. Thanks, however, to the silver suture and the courage of the operator, the operation was successfully performed and the patients cured. This, occurring in our good city, speaks volumes. For my own part, I think the evils resulting from severe lacerations are very great, and if anything I may say will direct more attention to the prevention of these evils I will be satisfied. I feel confident that the sum-total of the sorrow and misery arising from this cause vastly exceeds our conception. It is a recognized factor in the causation of subinvolution of the vagina and uterus, and I am persuaded its results are not limited to these organs, but that the tubes and varied ligaments share in the same mischief. It is a fruitful cause of retro-laxations of the uterus and prolapsus of bladder. Of all the marital misery and personal distress I need say nothing; these, of course, vary with the peculiarities of individual cases and the extent of the disease. I will not speak of the well-known preparation of the patient required, especially in extensive lacerations; you all know as to this and the after-treatment also. There is one remark I wish to make as to what is known as the perineal body. Some writers have made light of its existence, because its anatomy and relations are not sufficiently definite to merit, as they think, this appellation. That every uninjured perineum has such a body is unquestionable, and the restoration of this body is *the one* object of perineorrhaphy. An operation is successful or unsuccessful, according as to whether this end of the operation is or is not attained—without it the natural support of the pelvic viscera is impossible. Not only is there apt to be hernia of the anterior rectal wall, but prolapsus of both bladder and uterus—and this in the order I have given them. The best success heretofore has followed Emmet's operation. His conception of the trefoil character of the surfaces to be brought together are based upon a right conception of the anatomy of the parts. The perineal body being the central, and the lateral surfaces the outside leaves of the trefoil, each sulcus represents the lateral borders of the vagina and rectum. Perfect union of these surfaces leaves but little more to be desired. What remains to be attained is the object of what I now offer. In the first place, the loss of any tissue is

to be avoided, and sure union by first intention the desideratum to be attained. My operation is based upon the recognition of the immense value of the perineal body. I denude the surfaces to the fullest extent of the parts injured. This denudation is accomplished by the removal of the covering of the parts to be denuded—*i.e.*, the cicatricial surface in *one* piece. For this purpose the first incision is made at the upper part where the edge of the skin coalesces with the cicatricial surface—(the dotted line in sketch No. 1 shows this); the knife is entered at the highest point on the right side, and the incision brought down to the lowest part of the fourchette, when it is met by a similar incision on the left side. The lowest part of the angle is then seized with the forceps and carefully dissected upward, taking special care to remove the whole surface without incising the flap; this dissection is carried on till the surface represented by the original wound is uncovered. This flap, when raised with the hook, is seen in drawing No. 2. The next step is the introduction of the shield-sutures (and here I would say a word in favor of the catgut suture which I adopt). It is by far the best, as it gives the greatest possible extent of surface to surface—much greater than can be secured by the interrupted or any other suture. Two deep sutures usually suffice, and these—whether silver, silk, or catgut—are passed in and secured by clamp shot upon an ivory shield. The first suture should be inserted low down, and about three-quarters of an inch from the edge of the wound. It must be passed under the denuded surface so as not to appear, and brought out on the opposite side at a point corresponding to that of insertion. The second deep suture is similarly introduced higher up. The last deep suture should catch the flap, and the interrupted suture will do for this. The edges of the wound are coapted by horse-hair sutures, and the upper part of the flap and around on the right and left side are secured by catgut sutures; this leaves the united surfaces in the shape of the letter T. The vaginal surface is perfectly covered, and in no way can a drop of fluid enter the wound or interfere with union by first intention. There is very little pain, inasmuch as the deep shield-sutures allow of distention. Interrupted sutures should not be used. Where the rupture extends into the rectum, the flaps are carefully brought together by a running catgut suture, and the operation completed as in this case. The objection to all other