

Primaries brownish-black, with four or five large white or yellowish-white spots, and one or two small ones.

Secondaries light yellow with a brown spot near the anal angle.

Under surface of primaries with markings as above but much paler and overcast with yellow. Secondaries deeper in color than above.

Abdomen pale yellow with the tip of a deeper color, and a dorsal line of black. Legs yellow, the fore and middle pairs edged with black.

Length of body 6 lines. Wings expand 17-18.

Larva undescribed.

Hab.—Near West Flamboro'. Captured at midday in August. (Mr. Bethune.)

*H. interrupto-marginata*!—Beauv. Carolina, Harris. C. comma, Walker, C. B. M., 652. *Bornlix interrupto-marginata*. De Beauvois, Ins. Afriq. et Amer., p. 265, pl. 24.

Palpi deep ochre-yellow with black tips. Antennæ brown. Head pale orange-yellow. Thorax yellowish or pale buff, with a wide central black stripe, and a black spot on each side at base of primaries.

Primaries pale buff, with a black stripe along the costa not reaching the apex; a broad stripe of the same color along the inner margin, widening near the tip and sending from near the inner angle towards the hind end of the disk a hooked demi-band; hind margin with an incomplete black band, widest in the middle.

Secondaries bright orange-yellow with a black spot not far from the anal angle.

Under surface deep yellow throughout, with the markings on primaries excepting the demi-band scarcely perceptible.

Abdomen orange-yellow, with a dorsal band black. Legs deep yellow; the fore and middle pairs edged anteriorly with black.

Length of body 6-8 lines. Wings expand 16-20 lines.

Larva undescribed.

Hab.—London, rare; St. Catherines (Mr. Beadle); Port Stanley (Mr. Edwards); Wisconsin, and Virginia.