

Var. OCCIDENTALE, Trelease.

Lake Okanagan and Burrard Inlet, B.C., and common on Vancouver Island. "Sometimes comes too near *E. Franciscanum*, but differs in its usually smaller flowers less corymbosely clustered and more acute at base, and in its shorter glandular pubescence."

(15.) *E. GLANDULOSUM*, Lehm.

In damp places at an altitude of 5,000 feet at Warm Springs, Kootanie Lake, B.C. (*John Macoun*.)

(16.) *E. BREVISTYLUM*, Barbey.

Specimens from mountains south of Tulameen River B.C. (*Dawson*), have been doubtfully referred here by Dr. Trelease.

(17.) *E. HALLEANUM*, Hausskn.

Collected by Prof. Macoun in 1887 on Cedar Hill, Vancouver Island, and in 1893 at Esquimault, V. I.

(18.) *E. DRUMMONDII*, Hausskn.

Young specimens from Stewart's Lake, B. C., (*Macoun*) with leaves in whorls of 3, have been doubtfully referred here by Dr. Trelease.

(19.) *E. LEPTOCARPUM*, Hausskn.

A span or less high, glabrous except for some incurved pubescence on the stem; leaves less than 20 mm. long, broadly lanceolate, sparingly low-toothed, tapering from near the middle to the obtuse or subacute apex and winged petiole; flowers abundant for the size of the plant; calyx-tube narrow; petals about 3 mm. long, rosy; capsules 20 mm., on very slender peduncles of nearly equal length; seeds nearly ellipsoidal, shortly hyaline beaked, .25 × .75 mm.; coma at length cinnamon-colored.

Var. MACOUNII, Trelease.

Less branched, crisp-pubescent in lines, the same pubescence more or less abundant also on the flowers and capsules; leaves more ovate; seeds 1 mm. long; coma paler.