

4th. That the condition of the skin is similar to the condition of the mucous membrane in scarlatina.

In support of the first proposition, the microscopic examinations of the mucous membranes of the œsophagus, stomach and intestines were detailed in ten cases of death from scarlatina during the first week of illness, and in six cases who died in the second and third week of the fever. The first effects of the scarlatinal poison upon the mucous membrane of the stomach were shown to be the congestion of the bloodvessels and the stripping of the epithelium from the tubes and the surface of the organ, and also the softening of the tissues. The tubes are greatly distended by granular and fatty matters, or by small cells intermixed with granules, and in some cases they are lined by a newly-formed membrane. Sometimes no normal cells can be distinguished; in other cases they are present, but are scattered irregularly. After the second or third week the tubes are found less distended than at an earlier period, and whilst their closed ends are still loaded with granular matters, which greatly obscure the gastric cells. These become more evident towards the surface of the mucous membrane. The cells at this period are sometimes very large, sometimes loaded with fat or coated with granules, and seem to have but little adhesion to their basement membrane, as they readily separate from the tubes, but adhere closely to each other. The effects of the inflammation upon the intestines seem, in slighter cases, to consist in the effusion of granular and fatty matters into the mucous membrane, but in more severe cases the tubes of Lieberkühn are obstructed by epithelial cells, whilst extravasations of blood take place in the villi, and these, with the rest of the mucous membrane, are loaded with small cells and granules. In one case the mucous membrane was entirely stripped of villi, excepting a few fragments which still remained, and the enlarged and prominent openings of the follicles of Lieberkühn gave its surface the appearance of a sieve. In some instances in which the pancreas has been examined, evidences of disease presented themselves.

The second proposition was stated to be more difficult of proof, inasmuch as vomiting usually occurs only in the first stage, and the author had no opportunity of examining the vomited matters at this period of the disease. In one case, in which vomiting took place in the third week, fibrinous casts of the stomach tubes were discovered, and inflammation of the mucous membrane was proved to have existed by *post-mortem* examination. The chief reason upon which the opinion that desquamation of the epithelium occurs was founded, was from the microscopic examination of the stomachs of those who had died of this disease. The contents in recent cases consisted of pieces of fine membrane, of cells,