

Expectorants and Respiratory Sedatives for Horses—continued.

- ℞ P. fol. aconiti, ℥vi.
 P. digitalis, ℥iii.
 Arsenic. alb., grs. iv.
 P. anisi fruct., ℥ss.

Div. in pulv. vi. Sig. Give one every night in the food.
 (Useful in chronic cough.)

- ℞ Liq. arsenicalis, P.B., ℥iij.
 Tr. nucis vom., ℥iii.
 Tr. camphor. co., ℥iii.
 Aqua ad O.i.

F. m. Sig. Give two wineglassfuls twice a day after feeding.
 (Useful in coughs depending on gastric derangements.)

Dogs.

- ℞ Tr. belladonnæ, ℥ss.
 Syr. scillæ, ℥ss.
 Tr. camphor. co., ℥i.
 Aqua ad ℥vi.

F. m. Sig. Give two teaspoonfuls three times a

COUNTER-IRRITANTS.

- ℞ Pulv. cantharides, ℥xx.
 Ol. tereb. rect., ℥xii.
 Acid. acetic. fort., ℥ix.

Mix for twenty-four hours, then add 2½ lb. each of lanoline and vaseline, melted at a temperature of 200° Fahr., and stir well till thoroughly mixed. Keep in an air-tight jar.

Sig.: Ung. Cantharides Co. Fly b'ister.

- ℞ Hyd. biniod., ℥ii.
 Lanolini, ℥xii.
 Vaselini, ℥iv.

Ft. ung. Sig.: Ung. Hyd. Biniod.

(Useful for the treatment of ossific affections of joints, etc.)