

Many larvæ of some Cerambycide continued to work on under the bark ; late in the fall I observed the most of these had penetrated the wood, but some remained under the bark till April and May of the next year (1885). The most of the beetles appeared during the first two weeks of June, though individuals occurred occasionally till September. A few larvæ were still found at work, but by October they, likewise, had bored into the wood and appeared as beetles the next June (1886). The normal period of metamorphosis is therefore three years, but in individuals it may be retarded to four or more years.

At the present writing (June 5th) these beetles are issuing in great numbers from a barrel of hickory limbs obtained in April, 1885, from a tree deadened in January, 1884, thus verifying the first observation.

How the larvæ get under the bark could not be ascertained. When first examined, in April, they were from 4 to 5 m.m. long ; they ate the wood under the bark, following its grain, and packed their burrow solidly with their dust. Their growth and progress were both slow, for by the next April they had scarcely more than doubled in length, and had not traveled more than from four to six inches during the year ; but after July they developed an enormous appetite, and consumed the wood for at least an inch in length, and often entirely around the limb, ejecting their castings through holes made in the bark. When full fed, they bore obliquely an oval hole into the wood, penetrating it from four to ten inches. The larva then packs the opening with fine castings and enlarges a couple of inches of the interior of the burrow by gnawing off its sides a quantity of coarse fibre, in which it lies, after turning its head to the entrance. When about to become a pupa (I witnessed the process), the skin ruptures on the dorsum of three or four segments next the head ; the head of the pupa appears, and after about half an hour's wriggling the whole body is divested of its covering. To the observer the pupa appears to crawl out of the skin, but in fact the skin with the large mandibles is forced backwards by the alternate extension and contraction of the segments, assisted materially by the fibre that surrounds it. After its soft body hardens, the same movements free it from the fibre, some being shoved in advance of the head, and some posteriorly, the exuviae being often found at the distal end of the hole.

The time spent in the pupal state is indefinite, and does not seem to concern greatly the time of the appearance of the beetle. Sticks split open at different periods from December till March contained larvæ and