

## HEADACHE POWDERS.

Directions for use as found on the label.	Remarks.	RESULTS OF ANALYSIS.				No. of Sample.	Remarks and Opinion of the Chief Analyst.
		Weight of Contents of one powder, grains.	Acetanilide—Anti-febrin, grains.	Phenacetin, grains.	Phenazone—Antipyrim, grains.		

J. C. FERGUSON, INSPECTOR.—Continued.

Dose for headache.—One powder repeated in 30 minutes if necessary. At least three hours should elapse before another dose is taken. A seidlitz powder or other mild laxative should be taken with the first dose. Care should be taken in administering to children. Reduce the dose according to age. Dose for sleeplessness, one powder on retiring. The powders may be taken dry on the tongue followed by a little water.	Powder No. 1.....	11.6	4.5	None.	None.	45440	Acetanilide declared up to 4½ grains. One powder in excess of this amount. Meets requirements of P. or P. Act.
	" " 2.....	13.5	5.2	"	"		
Directions.—Place powder on the tongue and swallow with a draught of water. If not relieved in an hour another powder may be taken.	" " 1.....	14.3	none.	Less than 5 grains	none.	45441	Meets requirements of P. or P. Act.
	" " 2.....	12.9	"	"	"		
Directions.—Take one powder in a little water, and repeat in half an hour if not relieved. Take as early in the attack as possible.	Powder No. 1.....	10.3	2.1	none.	"	45442	Curative properties claimed. Meets requirements of P. or P. Act.
	" " 2.....	10.8	2.2	"	"		
Dose for Headache.—One powder repeated in thirty minutes if necessary, at least three hours should elapse before another dose is	Powder No. 1.....	10.9	4.1	"	"	45443	Acetanilide declared. Meets requirements of P. or P. Act.
	" " 2.....	11.8	4.6	"	"		