

many spatulate, linear, acute, entire leaves, and stoutish ascending, sparsely leafy monocephalous peduncles; the younger foliage canescently strigose, the older glabrate; heads large, hemispherical; bracts of the somewhat hoary-tomentose involucre subequal, in two series; rays about 50, rather broad, purple.

Summit of Sheep Mountain, Waterton Lake, Rocky Mountains, alt., 7,500 ft., July 31st, 1895. Herb. No. 10,858. (*John Macoun.*) Distributed as *Erigeron ochroleucus*.

ERIGERON KINDBERGI, Greene, Pittonia, Vol. III., p. 165.

Stems several, erect, from a perennial root, 6 inches high, pilose-pubescent, apparently flaccid and not conspicuously angled; lowest leaves oblanceolate, entire, acute, wholly glabrous and, in no degree, ciliate; the cauline narrowly linear, elongated, sessile by an abruptly dilated base; heads mostly solitary, small, the involucre barely three lines high; bracts very unequal, all narrowly linear and rather abruptly acute, glabrous and glandless, except at the pilose-pubescent base; rays very numerous, narrow and short; pappus scanty for the *E. acris* group, and not at all accrescent, dull-white, unchanged in age.

Meadows on the plateau east of Stump Lake, B.C., July 14th, 1891. Herb. No. 7,793. (*Jas. McEroy.*) Named for Dr. Nils Conrad Kindberg, by whom some of the characters were indicated in 1892.

ERIGERON JUCUNDUS, Greene, Pittonia, Vol. III., p. 165.

*E. acris*, Macoun, Cat. Can. Plants, Vol. I., p. 547 in part.

*E. acris*, var. *Dræbachensis*, Macoun, Cat. Can. Plants, Vol. I., p. 547 in part.

*E. alpinus*, Macoun, Cat. Can. Plants, Vol. I., p. 234 in part.

Perennial, 2 to 10 inches high, the several stems monocephalous, or in larger plants with several and corymbose-