

to the tip, which is bluntly pointed but not clavate; black, very faintly annulated with pale on a few basal segments; maxillary palpi short, brown, or blackish brown, base of cylindrical joints pale, last joint very little incurved, cut at tip, as long as 3rd and 4th together, 3rd a little longer than 4th, thicker on tip.

Labial palpi a little longer, basal joint pale; second longer, enlarged to tip, a little incurved; last joint longer, thicker to middle, above depressed, cylindrical, tip blunt; both joints blackish, pale on articulation.

Prothorax short, before the middle a transversal flat furrow, front margin slightly rounded; black with three yellow dots anteriorly and a posterior stripe on each side; with some white hairs, intermixed with black ones; mesothorax dull brownish gray, with a few scattered white hairs, two yellow dots anteriorly and four in a transversal series in the middle; the conical suture ending in the posterior margin pale yellow with a middle dagger-shaped black line, and on each side a shorter black line; in front of it two globular black shining elevations, which are approximate and like two ocelli; metathorax similar with some yellow spots.

Abdomen (male) longer than the wings, very slender, about cylindrical, blackish hirsute; brown, shining, darker below and at the apex; segments 2 to 4 with two pale dorsal longitudinal lines, which are sometimes partly confluent; the two following segments with two pale spots in middle; appendages brown, clothed densely with black hairs, straight, the base triangularly dilated to reach the dorsum of the segment; shorter than the last segment. Abdomen (female) much shorter than the wings, less slender, apical half thicker; color similar, but the long pale dorsal lines represented only by two middle and two apical spots; genital parts in the last segment with many strong black spines; upper part divided in two pale tubercles; below with two short brown appendages.

Legs short, pale, with white hairs, intermixed with a few black ones, principally at tip, densely sprinkled with black, the femurs sometimes nearly blackish; tip of tibia black; tarsi with apex of the two basal joints, the two following entirely, and tip of fifth, black; claws long, incurved, brown; spurs wanting.

Wings short, broadest before the bluntly-pointed apex; hyaline with white shades, a little fumose, the anteriors closely sprinkled with fuscous; venation dense, veins fuscous interrupted with white; around the transversals after the mediana and after the 4th vein, brown shades, sometimes forming brown streaks on the disk and near the hind margin; two series