

nearly purple, and has an orange tuft; venter with light yellow scales laterally, and an irregular median dark blue stripe extending the whole length of the abdomen.

Legs: Coxæ and trochanters dark, with brilliant light yellow scales; femora of fore legs light-scaled at base, with dark iridescent scales on the dorsal aspect for the distal two-thirds of its length; femora of mid-legs dark; femora of hind legs light for the proximal two-thirds, all a yellowish white ventrally; tibiæ of fore legs dark dorsally and light ventrally, except the very base and apex, which are a brilliant golden yellow; tibiæ of mid-legs mostly golden yellow, darker near the base, and a band of dark scales at the apex; tibiæ of hind legs dark blue, with green iridescence; metatarsi of fore legs nearly white, except a small basal spot of dark blue on the ventral aspect; of mid-leg nearly white, except a rather large dark spot on the dorsal surface; of hind legs dark blue, except a light basal band; 1st tarsal of fore leg is white, except an apical dark band, all the remaining joints dark; 1st tarsal on mid-leg and all the following are light yellow to white, except the last, which has a brown tip; 1st tarsal of hind legs is light, with a tiny dark apex, all the rest of the joints dark. Ungues simple and equal. In some lights there is a very narrow apical band of violet on all the femora, and on the mid and hind tibiæ reddish.

Wing clear reddish brown, sparsely scaled with brown, broadly truncate scales; 1st submarginal cell very short, very little more than half the length of 2nd posterior, and much narrower; mid and posterior cross-veins meet, and are about the same length; supernumerary cross-vein about half the length of mid, and three times its own length exterior to mid. Halteres orange brown.

There is a strong fold near fifth long vein, so that it makes a curvature in the margin of the wing.

Length, 12 + mm.

Habitat, Margosatubig, Mindanao, Philippine Islands.

Taken June and July.

Described from five specimens sent by Dr. H Newton Kierulf, Cont. Surg. U. S. Army, and evidently lies near *speciosus*, Skuse, and *Marshalli*, Theob., and may easily be the female of a species in which only the male has been known, but at all events it is sufficiently unlike such descriptions as I have been able to find to warrant my assuming it to be new.

(To be continued.)