

*EUDULE HYALINA*, n. sp.

This insect is of the size and shape of *E. unicolor*, Robs., from Texas, but, instead of being of its colour, is of the light straw colour of *E. mendica*, Wlk. It is apparently more hyaline than its congeners, and is very frail in its appearance. The wings are evenly unicolorous.

Senator, Ariz ; from Dr. Kunze. Taken September 9th, 1896.

*TEPHROCLYSTIS BOREALIS*, n. sp.

Expands 16 mm. Palpi fuscous, slightly ochreous ; front thorax and abdomen smooth silky fuscous, with a bluish tint. Fore wings fuscous, with cross lines fine, faint, the basal rounded, wavy, geminate, the middle wavy ; the outer heaviest towards and at costa, starting straight out at costa to vein 8 just beyond cell, following it towards base the distance of the angle from costa, then turning at an angle, almost a right angle, through discal spot straight, or nearly so, to inner margin. Hind wings fuscous, even, with wavy lines very faintly showing. Beneath fuscous, darker at costa on fore wings, and with hind wings darkened and lightened into 6 or 7 rounded parallel shadings.

Winnipeg, Manitoba ; from Mr. Hanham.

*TEPHROCLYSTIS LATIPENNIS*, n. sp.

Expands 16-18 mm. Palpi long, porrect, heavily scaled, black and fuscous mixed ; front somewhat tufted, fuscous or blackish gray ; thorax fuscous gray ; abdomen fuscous-gray to gray, segments sometimes tinged with brownish. Fore wings broad, gray to fuscous-gray, overlaid more or less with blackish, forming an extra basal, and marginal broad darker bands ; cross lines light, geminate, wavy ; one basal, the second within the prominent black discal spots, the outer extradiscal, the outer considerably bent below cell and often lined with black within ; a faint submarginal zigzag line, a marginal row of black dashes separated at end of veins. Hind wings broad, rounded, of same colour as fore wings, the basal and middle lines slightly indicated, the outer geminate, quite distinct, bent considerably at middle, wavy, margin darker ; a marginal black line ; discal points black, distinct. Beneath as above, the fore wings with markings more faint, the hind wings very nearly as above.

Quebec, Canada ; from Mr. Hanham.

*TEPHROCLYSTIS SUBCOLORATA*, n. sp.

Expands 20-23 mm. Palpi and head clear blackish ; thorax blackish-gray ; abdomen blackish-fuscous. Fore wings gray, overlaid with blackish ; the wings narrow extended, the lines quite distinct ; basal line